

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number  
WO 02/46227 A2(51) International Patent Classification: C07K 14/605,  
14/765, C12N 15/62, A61K 38/38, 38/26, C07K 19/00,  
A61P 3/00

(21) International Application Number: PCT/US01/43165

(22) International Filing Date:  
29 November 2001 (29.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(36) Priority Data:  
60/251,954 7 December 2000 (07.12.2000) US(71) Applicant (for all designated States except US): **ELI  
LILLY AND COMPANY** [US/US]; Lilly Corporate  
Center, Indianapolis, IN 46285 (US).(72) Inventors; and  
(73) Inventors/Applicants (for US only): **GLAESNER,  
Wolfgang** [DE/US]; 7512 Fieldstone Court, Indianapolis,  
IN 46254 (US); **MICANOVIC, Radmila** [US/US];  
7126 White Oak Trail, Indianapolis, IN 46236 (US);  
**TSCHANG, Sheng-Hung, Rainbow** [US/US]; 4963  
Riley Mews, Carmel, IN 46033 (US).(74) Agents: **STEWART, Mark, J. et al.**; Eli Lilly and Com-  
pany, Lilly Corporate Center, Drop Code 1104, Indianapo-  
lis, IN 46285 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU  
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DK,  
DK (utility model), DM, DZ, EC, EE, ES, FI (utility model),  
ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,  
NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility  
model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,  
YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

## Declarations under Rule 4.17:

--- as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(iii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,  
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,  
MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE,  
SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU,  
ZA, ZW. ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL,  
SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG,  
KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CI,  
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,  
TR). OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).--- as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(ii)) for the following desig-  
nations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY,  
BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC,  
EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN,  
IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,  
MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL,  
PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,  
UA, UG, UZ, VN, YU, ZA, ZW. ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

--- without international search reports and to be republished  
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the  
"Guidance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: GLP-1 FUSION PROTEINS

(57) Abstract: The present invention relates to glucagon-like-1 compounds fused to proteins that have the effect of extending the in  
vivo half-life of the peptides. These fusion proteins can be used to treat non-insulin dependent diabetes mellitus as well as a variety  
of other conditions.

WO 02/46227 A2

- 1 -

## GLP-1 FUSION PROTEINS

The present invention relates to glucagon-like peptides including analogs and derivatives thereof fused to proteins that have the effect of extending the in vivo half-life of the peptides. These fusion proteins can be used to treat non-insulin dependent diabetes mellitus as well as a variety of other conditions.

Glucagon-Like Peptide 1 (GLP-1) is a 37 amino acid peptide that is secreted by the L-cells of the intestine in response to food ingestion. It has been found to stimulate insulin secretion (insulinotropic action), thereby causing glucose uptake by cells and decreased serum glucose levels [see, e.g., Mojsov, S., (1992) *Int. J. Peptide Protein Research*, 40:333-343]. However, GLP-1 is poorly active. A subsequent endogenous cleavage between the 6<sup>th</sup> and 7<sup>th</sup> position produces a more potent biologically active GLP-1(7-37)OH peptide. Numerous GLP-1 analogs and derivatives are known and are referred to herein as "GLP-1 compounds." These GLP-1 analogs include the Exendins which are peptides found in the venom of the GILA-monster. The Exendins have sequence homology to native GLP-1 and can bind the GLP-1 receptor and initiate the signal transduction cascade responsible for the numerous activities that have been attributed to GLP-1(7-37)OH.

GLP-1 compounds have a variety of physiologically significant activities. For example, GLP-1 has been shown to stimulate insulin release, lower glucagon secretion, inhibit gastric emptying, and enhance glucose utilization. [Nauck, M.A., et al. (1993) *Diabetologia* 36:741-744; Gutniak, M., et al. (1992) *New England J. of Med.* 326:1316-1322; Nauck, M.A., et al., (1993) *J. Clin. Invest.* 91:301-307].

- 2 -

GLP-1 shows the greatest promise as a treatment for non-insulin dependent diabetes mellitus (NIDDM). There are numerous oral drugs on the market to treat the insulin resistance associated with NIDDM. As the disease progresses, however, patients must move to treatments that stimulate the release of insulin and eventually to treatments that involve injections of insulin. Current drugs which stimulate the release of insulin, however, can also cause hypoglycemia as can the actual administration of insulin. GLP-1 activity, however, is controlled by blood glucose levels. When levels drop to a certain threshold level, GLP-1 is not active. Thus, there is no risk of hypoglycemia associated with treatment involving GLP-1.

However, the usefulness of therapy involving GLP-1 peptides has been limited by their fast clearance and short half-lives. For example, GLP-1(7-37) has a serum half-life of only 3 to 5 minutes. GLP-1(7-36) amide has a time action of about 50 minutes when administered subcutaneously. Even analogs and derivatives that are resistant to endogenous protease cleavage, do not have half-lives long enough to avoid repeated administrations over a 24 hour period. Fast clearance of a therapeutic agent is inconvenient in cases where it is desired to maintain a high blood level of the agent over a prolonged period of time since repeated administrations will then be necessary. Furthermore, a long-acting compound is particularly important for diabetic patients whose past treatment regimen has involved taking only oral medication. These patients often have an extremely difficult time transitioning to a regimen that involves multiple injections of medication.

The present invention overcomes the problems associated with delivering a compound that has a short plasma half-life. The compounds of the present invention encompass GLP-1 compounds fused to another protein with a long circulating

- 3 -

half-life such as the Fc portion of an immunoglobulin or albumin.

Generally, small therapeutic peptides are difficult to manipulate because even slight changes in their structure can affect stability and/or biological activity. This has been especially true for GLP-1 compounds currently in development. For example, GLP-1(7-37)OH has a tendency to undergo a conformational change from a primarily alpha helix structure to a primarily beta sheet structure. This beta sheet form results in aggregated material that is thought to be inactive. It was, therefore, surprising that biologically active GLP-1 fusion proteins with increased half-lives could be developed. This was especially unexpected given the difficulty of working with GLP-1(7-37)OH alone and the large size of the fusion partner relative to the small GLP-1 peptide attached.

Compounds of the present invention include heterologous fusion proteins comprising a first polypeptide with a N-terminus and a C-terminus fused to a second polypeptide with a N-terminus and a C-terminus wherein the first polypeptide is a GLP-1 compound and the second polypeptide is selected from the group consisting of

- a) human albumin;
- b) human albumin analogs; and
- c) fragments of human albumin,

and wherein the C-terminus of the first polypeptide is fused to the N-terminus of the second polypeptide.

Compounds of the present invention also include a heterologous fusion protein comprising a first polypeptide with a N-terminus and a C-terminus fused to a second polypeptide with a N-terminus and a C-terminus wherein the first polypeptide is a GLP-1 compound and the second polypeptide is selected from the group consisting of

- a) human albumin;



- 4 -

- b) human albumin analogs; and
- c) fragments of human albumin,

and wherein the C-terminus of the first polypeptide is fused to the N-terminus of the second polypeptide via a peptide linker. It is preferred that the peptide linker is selected from the group consisting of:

- a) a glycine rich peptide;
- b) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>n</sub>, where n is 1, 2, 3, 4, 5 or 6; and
- 10 c) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>3</sub>.

Additional compounds of the present invention include a heterologous fusion protein comprising a first polypeptide with a N-terminus and a C-terminus fused to a second polypeptide with a N-terminus and a C-terminus wherein the first polypeptide is a GLP-1 compound and the second polypeptide is selected from the group consisting of

- a) the Fc portion of an immunoglobulin;
- b) an analog of the Fc portion of an immunoglobulin;
- 20 and

c) fragments of the Fc portion of an immunoglobulin, and wherein the C-terminus of the first polypeptide is fused to the N-terminus of the second polypeptide. The GLP-1 compound may be fused to the second polypeptide via a peptide linker. It is preferable that the peptide linker is selected from the group consisting of:

- a) a glycine rich peptide;
- b) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>n</sub>, where n is 1, 2, 3, 4, 5 or 6; and
- 30 c) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>3</sub>.

It is generally preferred that the GLP-1 compound that is part of the heterologous fusion protein have no more than 6 amino acids that are different from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4.

- 35 It is even more preferred that the GLP-1 compound have no

- 5 -

more than 5 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4. It is most preferred that the GLP-1 compound have no more than 4, 3, or 2 amino acids that differ from the

5 corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4. Preferably, a GLP-1 compound that is part of the heterologous fusion protein has glycine or valine at position 8.

The present invention also includes polynucleotides  
10 encoding the heterologous fusion protein described herein, vectors comprising these polynucleotides and host cells transfected or transformed with the vectors described herein. Also included is a process for producing a heterologous fusion protein comprising the steps of  
15 transcribing and translating a polynucleotide described herein under conditions wherein the heterologous fusion protein is expressed in detectable amounts.

The present invention also encompasses a method for normalizing blood glucose levels in a mammal in need thereof  
20 comprising the administration of a therapeutically effective amount of a heterologous fusion protein described herein.

The invention is further illustrated with reference to the following drawings:

25 Figure 1: IgG1 Fc amino acid sequence encompassing the hinge region, CH2 and CH3 domains.

Figure 2: Human serum albumin amino acid sequence.

Figure 3: A. SDS-PAGE gel and immunoblot of same gel illustrating the molecular weight of IgG1-Fc and GLP-1-Fc  
30 fusion proteins (Lane 1, MW standards; Lane 2, Purified Fc; lane 3, Mock transfected media; Lane 4, Val<sup>8</sup>-GLP-1-Fc; Lane 5, Exendin-4-Fc) B. SDS-PAGE gel and immunoblot of same gel illustrating the molecular weight of human HSA and GLP-1-HSA  
35 fusion proteins (Lane 1, MW standards; Lane 2, Purified HSA; lane 3, Mock transfected media; Lane 4, Val<sup>8</sup>-GLP-1-HSA; Lane

- 6 -

5, Val<sup>8</sup>-GLP-1-[Gly-Gly-Gly-Gly-Ser]<sub>3</sub>-HSA; Lane 6, Exendin-4-HSA; Lane 7, Exendin-4-[Gly-Gly-Gly-Gly-Ser]<sub>3</sub>-HSA).

Figure 4: SDS-PAGE gel of purified Fc, albumin, and GLP-1 fusion proteins (Lane 1, MW standards; Lane 2, purified Fc; Lane 3, Val<sup>8</sup>-GLP-1-Fc; Lane 4, Exendin-4-Fc; Lane 5, MW standard; Lane 6, Val<sup>8</sup>-GLP-1-HSA; Lane 7, Exendin-4-HSA; Lane 8, Exendin-4-[Gly-Gly-Gly-Gly-Ser]<sub>3</sub>-HSA).

Figure 5: Expression cloning vector containing the Fc regions illustrated in figure 1.

Figure 6: Expression cloning vector containing the albumin sequence illustrated in figure 2.

Figure 7: Expression cloning vector containing DNA encoding a 15 amino acid linker fused in frame and 5' of the albumin sequence illustrated in figure 2.

Figure 8: In vitro dose response activity of GLP-1 fusion proteins.

Figure 9: Pharmacokinetics of GLP-1 Fc and HSA fusion proteins.

Figure 10: Glucodynamic response to Exendin-Fc in two normal fasted dogs.

Figure 11: Insulinotropic response to Exendin-Fc in two normal fasted dogs.

Figure 12: DNA sequence encoding a human IgG1 Fc region.

Figure 13: DNA sequence encoding a human albumin protein.

The heterologous fusion proteins of the present invention comprise a GLP-1 compound fused to human albumin, a human albumin analog, a human albumin fragment, the Fc portion of an immunoglobulin, an analog of the Fc portion of an immunoglobulin, or a fragment of the Fc portion of an immunoglobulin. The C-terminus of the GLP-1 compound may be fused directly, or fused via a peptide linker, to the N-

- 7 -

terminus of an albumin or Fc protein. These heterologous fusion proteins are biologically active and have an increased half-life compared to native GLP-1.

It is preferred that the GLP-1 compounds that make up  
5 part of the heterologous fusion protein encompass polypeptides having from about twenty-five to about thirty-nine naturally occurring or non-naturally occurring amino acids that have sufficient homology to native GLP-1(7-37)OH such that they exhibit insulinotropic activity by binding to  
10 the GLP-1 receptor on  $\beta$ -cells in the pancreas. A GLP-1 compound typically comprises a polypeptide having the amino acid sequence of GLP-1(7-37)OH, an analog of GLP-1 (7-37)OH, a fragment of GLP-1(7-37)OH or a fragment of a GLP-1(7-37)OH analog. GLP-1(7-37)OH has the amino acid sequence of SEQ ID  
15 NO: 1:

7	8	9	10	11	12	13	14	15	16	17
His	Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Val	Ser
18	19	20	21	22	23	24	25	26	27	28
Ser	Tyr	Leu	Glu	Gly	Gln	Ala	Ala	Lys	Glu	Phe
29	30	31	32	33	34	35	36	37		
Ile	Ala	Trp	Leu	Val	Lys	Gly	Arg	Gly		

(SEQ ID NO: 1)

By custom in the art, the amino terminus of GLP-1(7-  
25 37)OH has been assigned number residue 7 and the carboxy-terminus, number 37. The other amino acids in the polypeptide are numbered consecutively, as shown in SEQ ID NO: 1. For example, position 12 is phenylalanine and position 22 is glycine.

30 GLP-1 compounds also encompass "GLP-1 fragments." A GLP-1 fragment is a polypeptide obtained after truncation of one or more amino acids from the N-terminus and/or C-terminus of GLP-1(7-37)OH or an analog or derivative thereof. The nomenclature used to describe GLP-1(7-37)OH is  
35 also applicable to GLP-1 fragments. For example, GLP-1(9-

- 8 -

36)OH denotes a GLP-1 fragment obtained by truncating two amino acids from the N-terminus and one amino acid from the C-terminus. The amino acids in the fragment are denoted by the same number as the corresponding amino acid in GLP-1(7-37)OH. For example, the N-terminal glutamic acid in GLP-1(9-36)OH is at position 9; position 12 is occupied by phenylalanine; and position 22 is occupied by glycine, as in GLP-1(7-37)OH. For GLP-1(7-36)OH, the glycine at position 37 of GLP-1(7-37)OH is deleted.

GLP-1 compounds also include polypeptides in which one or more amino acids have been added to the N-terminus and/or C-terminus of GLP-1(7-37)OH, or fragments or analogs thereof. It is preferred that GLP-1 compounds of this type have up to about thirty-nine amino acids. The amino acids in the "extended" GLP-1 compound are denoted by the same number as the corresponding amino acid in GLP-1(7-37)OH. For example, the N-terminus amino acid of a GLP-1 compound obtained by adding two amino acids to the N-terminal of GLP-1(7-37)OH is at position 5; and the C-terminus amino acid of a GLP-1 compound obtained by adding one amino acid to the C-terminus of GLP-1(7-37)OH is at position 38. Thus, position 12 is occupied by phenylalanine and position 22 is occupied by glycine in both of these "extended" GLP-1 compounds, as in GLP-1(7-37)OH. Amino acids 1-6 of an extended GLP-1 compound are preferably the same as or a conservative substitution of the amino acid at the corresponding position of GLP-1(1-37)OH. Amino acids 38-45 of an extended GLP-1 compound are preferably the same as or a conservative substitution of the amino acid at the corresponding position of glucagon or Exendin-4.

GLP-1 compounds of the present invention encompass "GLP-1 analogs." A GLP-1 analog has sufficient homology to GLP-1(7-37)OH or a fragment of GLP-1(7-37)OH such that the analog has insulinotropic activity. Preferably, a GLP-1 analog has the amino acid sequence of GLP-1(7-37)OH or a

- 9 -

fragment thereof, modified so that from one, two, three, four or five amino acids differ from the amino acid in the corresponding position of GLP-1(7-37)OH or a fragment of GLP-1(7-37)OH. In the nomenclature used herein to  
5 designate GLP-1 compounds, the substituting amino acid and its position is indicated prior to the parent structure. For example, Glu<sup>22</sup>-GLP-1(7-37)OH designates a GLP-1 compound in which the glycine normally found at position 22 of GLP-1(7-37)OH has been replaced with glutamic acid; Val<sup>8</sup>-Glu<sup>22</sup>-  
10 GLP-1(7-37)OH designates a GLP-1 compound in which alanine normally found at position 8 and glycine normally found at position 22 of GLP-1(7-37)OH have been replaced with valine and glutamic acid, respectively.

GLP-1 compounds of the present invention also include  
15 "GLP-1 derivatives." A GLP-1 derivative is defined as a molecule having the amino acid sequence of GLP-1 or of a GLP-1 analog, but additionally having chemical modification of one or more of its amino acid side groups,  $\alpha$ -carbon atoms, terminal amino group, or terminal carboxylic acid  
20 group. A chemical modification includes, but is not limited to, adding chemical moieties, creating new bonds, and removing chemical moieties. Modifications at amino acid side groups include, without limitation, acylation of lysine  $\epsilon$ -amino groups, N-alkylation of arginine, histidine, or  
25 lysine, alkylation of glutamic or aspartic carboxylic acid groups, and deamidation of glutamine or asparagine. Modifications of the terminal amino group include, without limitation, the des-amino, N-lower alkyl, N-di-lower alkyl, and N-acyl modifications. Modifications of the terminal  
30 carboxy group include, without limitation, the amide, lower alkyl amide, dialkyl amide, and lower alkyl ester modifications. Lower alkyl is C<sub>1</sub>-C<sub>4</sub> alkyl. Furthermore, one or more side groups, or terminal groups, may be protected by protective groups known to the ordinarily-

- 10 -

skilled protein chemist. The  $\alpha$ -carbon of an amino acid may be mono- or dimethylated.

Any GLP-1 compound can be part of the heterologous fusion proteins of the present invention as long as the GLP-1 compound itself is able to bind and induce signaling through the GLP-1 receptor. GLP-1 receptor binding and signal transduction can be assessed using *in vitro* assays such as those described in EP 619,322 and U.S. Patent No. 5,120,712, respectively.

10 Numerous active GLP-1 fragments, analogs and derivatives are known in the art and any of these analogs and derivatives can also be part of the heterologous fusion proteins of the present invention. Some examples of novel GLP-1 analogs as well as GLP-1 analogs and derivatives known  
15 in the art are provided herein.

Some GLP-1 analogs and GLP-1 fragments known in the art include, for example, GLP-1(7-34) and GLP-1(7-35), GLP-1(7-36), Gln<sup>9</sup>-GLP-1(7-37), D-Gln<sup>9</sup>-GLP-1(7-37), Thr<sup>16</sup>-Lys<sup>18</sup>-GLP-1(7-37), and Lys<sup>18</sup>-GLP-1(7-37). GLP-1 analogs such as GLP-1(7-34) and GLP-1(7-35) are disclosed in U.S. Patent No.  
20 5,118,666. Biologically processed forms of GLP-1 which have insulinotropic properties, such as GLP-1(7-36) are also known. Other known biologically active GLP-1 compounds are disclosed in U.S. Patent No 5,977,071 to Hoffmann, et al.,  
25 U.S. Patent No. 5,545,618 to Buckley, et al., and Adelhorst, et al., *J. Biol. Chem.* 269:6275 (1994).

- 11 -

A preferred group of GLP-1 analogs is composed of  
GLP-1 analogs of formula I (SEQ ID NO: 2)

```

5           7   8   9   10  11  12  13  14  15  16  17
           His-Xaa-Xaa-Gly-Xaa-Phe-Thr-Xaa-Asp-Xaa-Xaa-
           18  19  20  21  22  23  24  25  26  27  28
           Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Phe-
           29  30  31  32  33  34  35  36  37  38  39
10      Ile-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-
           40  41  42  43  44  45
           Xaa-Xaa-Xaa-Xaa-Xaa-Xaa

```

Formula I (SEQ ID NO: 2)

15           wherein:

```

Xaa at position 8 is Ala, Gly, Ser, Thr, Leu, Ile, Val,
    Glu, Asp, or Lys;
Xaa at position 9 is Glu, Asp, or Lys;
Xaa at position 11 is Thr, Ala, Gly, Ser, Leu, Ile, Val,
20     Glu, Asp, or Lys;
Xaa at position 14 is Ser, Ala, Gly, Thr, Leu, Ile, Val,
    Glu, Asp, or Lys;
Xaa at position 16 is Val, Ala, Gly, Ser, Thr, Leu, Ile,
    Tyr, Glu, Asp, Trp, or Lys;
25 Xaa at position 17 is Ser, Ala, Gly, Thr, Leu, Ile, Val,
    Glu, Asp, or Lys;
Xaa at position 18 is Ser, Ala, Gly, Thr, Leu, Ile, Val,
    Glu, Asp, Trp, Tyr, or Lys;
Xaa at position 19 is Tyr, Phe, Trp, Glu, Asp, Gln, or Lys;
30 Xaa at position 20 is Leu, Ala, Gly, Ser, Thr, Ile, Val,
    Glu, Asp, Met, Trp, Tyr, or Lys;
Xaa at position 21 is Glu, Asp, or Lys;
Xaa at position 22 is Gly, Ala, Ser, Thr, Leu, Ile, Val,
    Glu, Asp, or Lys;
35 Xaa at position 23 is Gln, Asn, Arg, Glu, Asp, or Lys;

```



- 12 -

- Xaa at position 24 is Ala, Gly, Ser, Thr, Leu, Ile, Val,  
Arg, Glu, Asp, or Lys;  
Xaa at position 25 is Ala, Gly, Ser, Thr, Leu, Ile, Val,  
Glu, Asp, or Lys;  
5 Xaa at position 26 is Lys, Arg, Gln, Glu, Asp, or His;  
Xaa at position 27 is Leu, Glu, Asp, or Lys;  
Xaa at position 30 is Ala, Gly, Ser, Thr, Leu, Ile, Val,  
Glu, Asp, or Lys;  
Xaa at position 31 is Trp, Phe, Tyr, Glu, Asp, or Lys;  
10 Xaa at position 32 is Leu, Gly, Ala, Ser, Thr, Ile, Val,  
Glu, Asp, or Lys;  
Xaa at position 33 is Val, Gly, Ala, Ser, Thr, Leu, Ile,  
Glu, Asp, or Lys;  
Xaa at position 34 is Asn, Lys, Arg, Glu, Asp, or His;  
15 Xaa at position 35 is Gly, Ala, Ser, Thr, Leu, Ile, Val,  
Glu, Asp, or Lys;  
Xaa at position 36 is Gly, Arg, Lys, Glu, Asp, or His;  
Xaa at position 37 is Pro, Gly, Ala, Ser, Thr, Leu, Ile,  
Val, Glu, Asp, or Lys, or is deleted;  
20 Xaa at position 38 is Ser, Arg, Lys, Glu, Asp, or His, or is  
deleted;  
Xaa at position 39 is Ser, Arg, Lys, Glu, Asp, or His, or is  
deleted;  
Xaa at position 40 is Gly, Asp, Glu, or Lys, or is deleted;  
25 Xaa at position 41 is Ala, Phe, Trp, Tyr, Glu, Asp, or Lys,  
or is deleted;  
Xaa at position 42 is Ser, Pro, Lys, Glu, or Asp, or is  
deleted;  
Xaa at position 43 is Ser, Pro, Glu, Asp, or Lys, or is  
30 deleted;  
Xaa at position 44 is Gly, Pro, Glu, Asp, or Lys, or is  
deleted;  
and  
Xaa at position 45 is Ala, Ser, Val, Glu, Asp, or Lys, or is  
35 deleted;

- 13 -

provided that when the amino acid at position 37, 38, 39, 40, 41, 42, 43, or 44 is deleted, then each amino acid downstream of that amino acid is also deleted.

It is preferred that the GLP-1 compound of formula I contain less than six amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH or Exendin-4. It is more preferred that less than five amino acids differ from the corresponding amino acid in GLP-1(7-37)OH or Exendin-4. It is even more preferred that less than four amino acids differ from the corresponding amino acid in GLP-1(7-37)OH or Exendin-4.

GLP-1 compounds of the present invention include derivatives of formula I such as a C-1-6-ester, or amide, or C-1-6-alkylamide, or C-1-6-dialkylamide thereof. WO99/43706 describes derivatives of GLP-1 compounds of formula I and is incorporated by reference herein in its entirety. The compounds of formula I derivatized as described in WO99/43706 and underivatized are encompassed by the present invention.

Another preferred group of GLP-1 compounds is composed of GLP-1 analogs of formula II (SEQ ID NO: 3):

7 8 9 10 11 12 13 14 15 16 17  
Xaa-Xaa-Xaa-Gly-Xaa-Xaa-Thr-Ser-Asp-Xaa-Ser-  
18 19 20 21 22 23 24 25 26 27 28  
Xaa-Xaa-Leu-Glu-Gly-Xaa-Xaa-Ala-Xaa-Xaa-Phe-  
29 30 31 32 33 34 35 36 37  
Ile-Xaa-Xaa-Leu-Xaa-Xaa-Xaa-Xaa-R

Formula II (SEQ ID NO: 3)

30

wherein:

Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;

35

- 14 -

- Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;  
Xaa at position 9 is: Thr, Ser, Arg, Lys, Trp, Phe, Tyr,  
Glu, or His;  
Xaa at position 11 is: Asp, Glu, Arg, Thr, Ala, Lys, or His;  
5 Xaa at position 12 is: His, Trp, Phe, or Tyr;  
Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp,  
Val, Tyr, Glu, or Ala;  
Xaa at position 18 is: His, Pro, Asp, Glu, Arg, Ser, Ala, or  
Lys;  
10 Xaa at position 19 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or  
Cys;  
Xaa at position 23 is: His, Asp, Lys, Glu, Gln, or Arg;  
Xaa at position 24 is: Glu, Arg, Ala, or Lys;  
Xaa at position 26 is: Trp, Tyr, Phe, Asp, Lys, Glu, or His;  
15 Xaa at position 27 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or  
Lys;  
Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;  
Xaa at position 31 is: Asp, Glu, Ser, Thr, Arg, Trp, or Lys;  
Xaa at position 33 is: Asp, Arg, Val, Lys, Ala, Gly, or Glu;  
20 Xaa at position 34 is: Glu, Lys, or Asp;  
Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe,  
Asp, Gly, Pro, His, or Glu;  
Xaa at position 36 is: Thr, Ser, Asp, Trp, Tyr, Phe, Arg,  
Glu, or His;  
25 R at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr,  
Phe, His, Gly, Gly-Pro, or is deleted.

- 15 -

Another preferred group of GLP-1 compounds is composed of GLP-1 analogs of formula III (SEQ ID NO: 4):

	7	8	9	10	11	12	13	14	15	16	17
5	Xaa	Xaa	Glu	Gly	Xaa	Xaa	Thr	Ser	Asp	Xaa	Ser
	18	19	20	21	22	23	24	25	26	27	28
	Ser	Tyr	Leu	Glu	Xaa	Xaa	Xaa	Xaa	Lys	Xaa	Phe
	29	30	31	32	33	34	35	36	37		
	Ile	Xaa	Trp	Leu	Xaa	Xaa	Xaa	Xaa	R		
10	formula III (SEQ ID NO: 4)										

wherein:

- Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine,  
15 homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;
- Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;
- Xaa at position 11 is: Asp, Glu, Arg, Thr, Ala, Lys, or His;
- Xaa at position 12 is: His, Trp, Phe, or Tyr;
- 20 Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp, Val, Glu, or Ala;
- Xaa at position 22: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;
- Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;
- 25 Xaa at position 24 is: Glu, His, Ala, or Lys;
- Xaa at position 25 is: Asp, Lys, Glu, or His;
- Xaa at position 27 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys;
- Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;
- 30 Xaa at position 33 is: Asp, Arg, Val, Lys, Ala, Gly, or Glu;
- Xaa at position 34 is: Glu, Lys, or Asp;
- Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;
- Xaa at position 36 is: Arg, Glu, or His;

- 16 -

R at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

Another preferred group of GLP-1 compounds is composed  
5 of GLP-1 analogs of formula IV (SEQ ID NO: 5):

7 8 9 10 11 12 13 14 15 16 17  
Xaa-Xaa-Glu-Gly-Thr-Xaa-Thr-Ser-Asp-Xaa-Ser-  
18 19 20 21 22 23 24 25 26 27 28  
Ser-Tyr-Leu-Glu-Xaa-Xaa-Ala-Ala-Xaa-Glu-Phe-  
10 29 30 31 32 33 34 35 36 37  
Ile-Xaa-Trp-Leu-Val-Lys-Xaa-Arg-R  
formula IV (SEQ ID NO: 5)

wherein:

- 15 Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;
- Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, Met, or  
20 Thr;
- Xaa at position 12 is: His, Trp, Phe, or Tyr;
- Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp,  
Val, Glu, or Ala;
- Xaa at position 22 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or  
25 Cys;
- Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;
- Xaa at position 26 is: Asp, Lys, Glu, or His;
- Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;
- Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe,  
30 Asp, Gly, Pro, His, or Glu;
- R at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

- 17 -

Another preferred group of GLP-1 compounds is composed of GLP-1 analogs of formula V (SEQ ID NO: 6):

5                   7   8   9   10 11 12 13 14 15 16 17  
                  Xaa-Xaa-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-  
                  18 19 20 21 22 23 24 25 26 27 28  
                  Ser-Tyr-Leu-Glu-Xaa-Xaa-Xaa-Ala-Lys-Glu-Phe-  
                  29 30 31 32 33 34 35 36 37  
                  Ile-Xaa-Trp-Leu-Val-Lys-Gly-Arg-R  
10                               formula V (SEQ ID NO: 6)

wherein:

Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;

15                   Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;

Xaa at position 22 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;

Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;

20                   Xaa at position 24 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys;

Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;

R at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

25                   Preferred GLP-1 compounds of formula I, II, III, IV, and V comprise GLP-1 analogs or fragments of GLP-1 analogs wherein the analogs or fragments contain an amino acid other than alanine at position 8 (position 8 analogs). It is

30                   preferable that these position 8 analogs contain one or more additional changes at positions 9, 11, 12, 16, 18, 22, 23, 24, 26, 27, 30, 31, 33, 34, 35, 36, and 37 compared to the corresponding amino acid of native GLP-1(7-37)OH. It is

also preferable that these analogs have 6 or fewer changes

35                   compared to the corresponding amino acids in native GLP-1(7-

- 18 -

37)OH or GLP-1(7-36)OH. More preferred analogs have 5 or fewer changes compared to the corresponding amino acids in native GLP-1(7-37)OH or GLP-1(7-36)OH or have 4 or fewer changes compared to the corresponding amino acids in native GLP-1(7-37)OH or GLP-1(7-36)OH. It is even more preferable that these analogs have 3 or fewer changes compared to the corresponding amino acids in native GLP-1(7-37)OH or GLP-1(7-36)OH. It is most preferable that these analogs have 2 or fewer changes compared to the corresponding amino acids in native GLP-1(7-37)OH.

It has been found that the compounds of formula II, III, IV, and V have a reduced propensity to aggregate and generate insoluble forms. This is also important in the context of a fusion protein wherein the relatively small GLP-1 peptide must maintain an active conformation despite being fused to a much larger protein. Preferred GLP-1 compounds of formula II, III, IV, and V encompassed by the fusion proteins of the present invention comprise GLP-1 analogs or fragments of GLP-1 analogs in which glycine at position 22 and preferably alanine at position 8 have been replaced with another amino acid.

When position 22 is aspartic acid, glutamic acid, arginine or lysine, position 8 is preferably glycine, valine, leucine, isoleucine, serine, threonine or methionine and more preferably valine or glycine. When position 22 is a sulfonic acid such as cysteic acid, position 8 is preferably glycine, valine, leucine, isoleucine, serine, threonine or methionine and more preferably valine or glycine.

Other preferred GLP-1 compounds include GLP-1 analogs of formula IV (SEQ ID NO:5) wherein the analogs have the sequence of GLP-1(7-37)OH except that the amino acid at position 8 is preferably glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine and position 30 is glutamic

- 19 -

acid, aspartic acid, serine, or histidine and more preferably glutamatic acid.

Other preferred GLP-1 compounds include GLP-1 analogs of formula IV (SEQ ID NO:5) wherein the analogs have the  
5 sequence of GLP-1(7-37)OH except that the amino acid at position 8 is preferably glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine and position 37 is histidine, lysine, arginine, threonine, serine, glutamic acid, aspartic  
10 acid, tryptophan, tyrosine, phenylalanine and more preferably histidine.

Other preferred GLP-1 compounds include GLP-1 analogs of formula IV (SEQ ID NO:5) wherein the analogs have the sequence of GLP-1(7-37)OH except that the amino acid at  
15 position 8 is preferably glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine and position 22 is glutamic acid, lysine, aspartic acid, or arginine and more preferably glutamic acid or lysine and position 23 is lysine, arginine,  
20 glutamic acid, aspartic acid, and histidine and more preferably lysine or glutamic acid.

Other preferred GLP-1 compounds include GLP-1 analogs of formula V (SEQ ID NO:6) wherein the analogs have the sequence of GLP-1(7-37)OH except that the amino acid at  
25 position 8 is preferably glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine and position 22 is glutamic acid, lysine, aspartic acid, or arginine and more preferably glutamine acid or lysine and position 27 is alanine, lysine,  
30 arginine, tryptophan, tyrosine, phenylalanine, or histidine and more preferably alanine.

Other preferred GLP-1 compounds include GLP-1 analogs of formula II wherein the analogs have the sequence of GLP-



- 20 -

1(7-37)OH except that the amino acid at position 8 and one, two, or three amino acids selected from the group consisting of position 9, position 11, position 12, position 16, position 18, position 22, position 23, position 24, position 26, position 27, position 30, position 31, position 33, position 34, position 35, position 36, and position 37, differ from the amino acid at the corresponding position of native GLP-1(7-37)OH.

Other preferred GLP-1 compounds of formula II include:

Val<sup>8</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-GLP-1(7-37)OH, Glu<sup>22</sup>-GLP-1(7-37)OH, Asp<sup>22</sup>-GLP-1(7-37)OH, Arg<sup>22</sup>-GLP-1(7-37)OH, Lys<sup>22</sup>-GLP-1(7-37)OH, Cys<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Arg<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Cys<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Arg<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Cys<sup>22</sup>-GLP-1(7-37)OH, Glu<sup>22</sup>-GLP-1(7-36)OH, Asp<sup>22</sup>-GLP-1(7-36)OH, Arg<sup>22</sup>-GLP-1(7-36)OH, Lys<sup>22</sup>-GLP-1(7-36)OH, Cys<sup>22</sup>-GLP-1(7-36)OH, Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)OH, Val<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-36)OH, Val<sup>8</sup>-Arg<sup>22</sup>-GLP-1(7-36)OH, Val<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-36)OH, Val<sup>8</sup>-Cys<sup>22</sup>-GLP-1(7-36)OH, Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)OH, Gly<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-36)OH, Gly<sup>8</sup>-Arg<sup>22</sup>-GLP-1(7-36)OH, Gly<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-36)OH, Gly<sup>8</sup>-Cys<sup>22</sup>-GLP-1(7-36)OH, Lys<sup>23</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Lys<sup>23</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Lys<sup>23</sup>-GLP-1(7-37)OH, His<sup>24</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-His<sup>24</sup>-GLP-1(7-37)OH, Lys<sup>24</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Lys<sup>24</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Lys<sup>23</sup>-GLP-1(7-37)OH, Glu<sup>30</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-37)OH, Asp<sup>30</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Asp<sup>30</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Asp<sup>30</sup>-GLP-1(7-37)OH, Gln<sup>30</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Gln<sup>30</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Gln<sup>30</sup>-GLP-1(7-37)OH, Tyr<sup>30</sup>-

- 21 -

GLP-1(7-37)OH, Val<sup>8</sup>-Tyr<sup>30</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Tyr<sup>30</sup>-GLP-1(7-37)OH, Ser<sup>30</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Ser<sup>30</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Ser<sup>30</sup>-GLP-1(7-37)OH, His<sup>30</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-His<sup>30</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-His<sup>30</sup>-GLP-1(7-37)OH, Glu<sup>34</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>34</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Glu<sup>34</sup>-GLP-1(7-37)OH, Ala<sup>34</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Ala<sup>34</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Ala<sup>34</sup>-GLP-1(7-37)OH, Gly<sup>34</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Gly<sup>34</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Gly<sup>34</sup>-GLP-1(7-37)OH, Ala<sup>35</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Ala<sup>35</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Ala<sup>35</sup>-GLP-1(7-37)OH, Lys<sup>35</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Lys<sup>35</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Lys<sup>35</sup>-GLP-1(7-37)OH, His<sup>35</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-His<sup>35</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-His<sup>35</sup>-GLP-1(7-37)OH, Pro<sup>35</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Pro<sup>35</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Pro<sup>35</sup>-GLP-1(7-37)OH, Glu<sup>35</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>35</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Glu<sup>35</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Ala<sup>27</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>22</sup>-Lys<sup>23</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>22</sup>-Glu<sup>23</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>22</sup>-Ala<sup>27</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Gly<sup>34</sup>-Lys<sup>35</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>22</sup>-Ala<sup>27</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Glu<sup>22</sup>-Ala<sup>27</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Lys<sup>22</sup>-Glu<sup>23</sup>-GLP-1(7-37)OH, and Gly<sup>8</sup>-Lys<sup>22</sup>-Glu<sup>23</sup>-GLP-1(7-37)OH.

Another preferred group of GLP-1 analogs and derivatives for use in the present invention is composed of molecules of formula VI (SEQ ID NO: 7)

25 R1-X-Glu-Gly<sup>10</sup>-Thr-Phe-Thr-Ser-Asp<sup>15</sup>-Val-Ser-  
Ser-Tyr-Leu<sup>20</sup>-Y -Gly-Gln-Ala-Ala<sup>25</sup>-Lys- Z -  
Phe-Ile-Ala<sup>30</sup>-Trp-Leu-Val-Lys-Gly<sup>35</sup>-Arg-R2  
formula VI (SEQ ID NO:7)

- 22 -

wherein: R<sub>1</sub> is selected from the group consisting of L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine, alpha-fluoromethyl-histidine, and alpha-methyl-histidine; X is  
5 selected from the group consisting of Ala, Gly, Val, Thr, Ile, and alpha-methyl-Ala; Y is selected from the group consisting of Glu, Gln, Ala, Thr, Ser, and Gly; Z is selected from the group consisting of Glu, Gln, Ala, Thr, Ser, and Gly; and R<sub>2</sub> is Gly-OH.

- 10 Another preferred group of GLP-1 compounds for use in the present invention is disclosed in WO 91/11457, and consists essentially of GLP-1(7-34), GLP-1(7-35), GLP-1(7-36), or GLP-1(7-37), or the amide form thereof, and pharmaceutically-acceptable salts thereof, having at least  
15 one modification selected from the group consisting of:

- (a) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, phenylalanine, arginine, or D-lysine for lysine at position 26 and/or position 34; or  
20 substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, phenylalanine, lysine, or a D-arginine for arginine at position 36;

- (b) substitution of an oxidation-resistant amino  
25 acid for tryptophan at position 31;

- (c) substitution of at least one of: tyrosine for valine at position 16; lysine for serine at position 18; aspartic acid for glutamic acid at position 21; serine for glycine at position 22; arginine for glutamine at position  
30 23; arginine for alanine at position 24; and glutamine for lysine at position 26; and

- (d) substitution of at least one of: glycine, serine, or cysteine for alanine at position 8; aspartic acid, glycine, serine, cysteine, threonine, asparagine,  
35 glutamine, tyrosine, alanine, valine, isoleucine, leucine,

- 23 -

methionine, or phenylalanine for glutamic acid at position 9; serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine for glycine at position 10; and glutamic acid for aspartic acid at position 15; and

(e) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine, or the D- or N-acylated or alkylated form of histidine for histidine at position 7; wherein, in the substitutions is (a), (b), (d), and (e), the substituted amino acids can optionally be in the D-form and the amino acids substituted at position 7 can optionally be in the N-acylated or N-alkylated form.

Because the enzyme, dipeptidyl-peptidase IV (DPP IV), may be responsible for the observed rapid *in vivo* inactivation of administered GLP-1, [see, e.g., Mentlein, R., et al., *Eur. J. Biochem.*, 214:829-835 (1993)], GLP-1 analogs and derivatives that are protected from the activity of DPP IV in the context of a fusion protein are preferred, and fusion proteins wherein the GLP-1 compound is Gly<sup>8</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-GLP-1(7-37)OH,  $\alpha$ -methyl-Ala<sup>8</sup>-GLP-1(7-37)OH, or Gly<sup>8</sup>-Gln<sup>21</sup>-GLP-1(7-37)OH are more preferred.

Another preferred group of GLP-1 compounds for use in the present invention consists of the compounds of formula VII (SEQ ID NO: 8) claimed in U.S. Patent No. 5,512,549, which is expressly incorporated herein by reference.

R<sup>1</sup>-Ala-Glu-Gly<sup>10</sup>-  
Thr-Phe-Thr-Ser-Asp<sup>15</sup>-Val-Ser-Ser-Tyr-Leu<sup>20</sup>-  
Glu-Gly-Gln-Ala-Ala<sup>25</sup>-Xaa-Glu-Phe-Ile-Ala<sup>30</sup>-  
Trp-Leu-Val-Lys-Gly<sup>35</sup>-Arg-R<sup>3</sup>

|  
R<sup>2</sup>

formula VII (SEQ ID NO: 8)

35

- 24 -

wherein R<sup>1</sup> is selected from the group consisting of 4-imidazopropionyl, 4-imidazoacetyl, or 4-imidazo- $\alpha,\alpha$ -dimethyl-acetyl; R<sup>2</sup> is selected from the group consisting of C<sub>6</sub>-C<sub>10</sub> unbranched acyl, or is absent; R<sup>3</sup> is  
5 selected from the group consisting of Gly-OH or NH<sub>2</sub>; and, Xaa is Lys or Arg.

More preferred compounds of formula IV for use in the present invention are those in which Xaa is Arg and R<sup>2</sup> is C<sub>6</sub>-C<sub>10</sub> unbranched acyl. Even more preferred compounds of  
10 formula IV for use in the present invention are those in which Xaa is Arg, R<sup>2</sup> is C<sub>6</sub>-C<sub>10</sub> unbranched acyl, and R<sup>3</sup> is Gly-OH. Other highly-preferred compounds of formula IV for use in the present invention are those in which Xaa is Arg, R<sup>2</sup> is C<sub>6</sub>-C<sub>10</sub> unbranched acyl, R<sup>3</sup> is Gly-OH, and R<sup>1</sup> is 4-  
15 imidazopropionyl. An especially preferred compound of formula IV for use in the present invention is that in which Xaa is Arg, R<sup>2</sup> is C<sub>8</sub> unbranched acyl, R<sup>3</sup> is Gly-OH, and R<sup>1</sup> is 4-imidazopropionyl.

Preferably, the GLP-1 compounds comprise GLP-1 analogs  
20 wherein the backbone for such analogs or fragments contains an amino acid other than alanine at position 8 (position 8 analogs). The backbone may also include L-histidine, D-histidine, or modified forms of histidine such as desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine,  
25 homohistidine,  $\alpha$ -fluoromethyl-histidine, or  $\alpha$ -methyl-histidine at position 7. It is preferable that these position 8 analogs contain one or more additional changes at positions 12, 16, 18, 19, 20, 22, 25, 27, 30, 33, and 37  
30 37)OH. It is more preferable that these position 8 analogs contain one or more additional changes at positions 16, 18,

- 25 -

22, 25 and 33 compared to the corresponding amino acid of native GLP-1(7-37)OH.

In a preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 12 is selected from the group consisting of tryptophan or tyrosine. It is more preferred that in addition to the substitution at position 12, the amino acid at position 8 is substituted with glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine. It is even more preferred that in addition to the substitutions at position 12 and 8, the amino acid at position 22 is substituted with glutamic acid.

In another preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 16 is selected from the group consisting of tryptophan, isoleucine, leucine, phenylalanine, or tyrosine. It is more preferred that in addition to the substitution at position 16, the amino acid at position 8 is substituted with glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine. It is even more preferred that in addition to the substitutions at position 16 and 8, the amino acid at position 22 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 16 and 8, the amino acid at position 30 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 16 and 8, the amino acid at position 37 is substituted with histidine.

In another preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 18 is selected from the group consisting of tryptophan, tyrosine, phenylalanine, lysine, leucine, or isoleucine, preferably tryptophan, tyrosine, and isoleucine. It is more preferred

- 26 -

that in addition to the substitution at position 18, the amino acid at position 8 is substituted with glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine. It is even more preferred that in addition to the substitutions at position 18 and 8, the amino acid at position 22 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 18 and 8, the amino acid at position 30 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 18 and 8, the amino acid at position 37 is substituted with histidine

In another preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 19 is selected from the group consisting of tryptophan or phenylalanine, preferably tryptophan. It is more preferred that in addition to the substitution at position 19, the amino acid at position 8 is substituted with glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine. It is even more preferred that in addition to the substitutions at position 19 and 8, the amino acid at position 22 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 19 and 8, the amino acid at position 30 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 19 and 8, the amino acid at position 37 is substituted with histidine

In another preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 20 is phenylalanine, tyrosine, or tryptophan. It is more preferred that in addition to the substitution at position 20, the amino acid at position 8 is substituted with

- 27 -

glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine. It is even more preferred that in addition to the substitutions at position 20 and 8, the amino acid at position 22 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 20 and 8, the amino acid at position 30 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 20 and 8, the amino acid at position 37 is substituted with histidine

In another preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 25 is selected from the group consisting of valine, isoleucine, and leucine, preferably valine. It is more preferred that in addition to the substitution at position 25, the amino acid at position 8 is substituted with glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine. It is even more preferred that in addition to the substitutions at position 25 and 8, the amino acid at position 22 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 25 and 8, the amino acid at position 30 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 25 and 8, the amino acid at position 37 is substituted with histidine.

In another preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 27 is selected from the group consisting of isoleucine or alanine. It is more preferred that in addition to the substitution at position 27, the amino acid at position 8 is substituted with glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or



- 28 -

glycine. It is even more preferred that in addition to the substitutions at position 27 and 8, the amino acid at position 22 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 27 and 8, the amino acid at position 30 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 27 and 8, the amino acid at position 37 is substituted with histidine

In another preferred embodiment, the GLP-1 analog is GLP-1(7-37)OH wherein the amino acid at position 33 is isoleucine. It is more preferred that in addition to the substitution at position 33, the amino acid at position 8 is substituted with glycine, valine, leucine, isoleucine, serine, threonine, or methionine and more preferably valine or glycine. It is even more preferred that in addition to the substitutions at position 33 and 8, the amino acid at position 22 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 33 and 8, the amino acid at position 30 is substituted with glutamic acid. It is also preferred that in addition to the substitutions at positions 33 and 8, the amino acid at position 37 is substituted with histidine

The GLP-1 compounds have modifications at one or more of the following positions: 8, 12, 16, 18, 19, 20, 22, 25, 27, 30, 33, and 37. These GLP-1 compounds show increased potency compared with GLP-1(7-37)OH and comprise the amino acid sequence of formula IX (SEQ ID NO:12)

Xaa<sub>7</sub>-Xaa<sub>8</sub>-Glu-Gly-Thr-Xaa<sub>12</sub>-Thr-Ser-Asp-Xaa<sub>16</sub>-Ser-Xaa<sub>18</sub>-Xaa<sub>19</sub>-Xaa<sub>20</sub>-Glu-Xaa<sub>22</sub>-Gln-Ala-Xaa<sub>25</sub>-Lys-Xaa<sub>27</sub>-Phe-Ile-Xaa<sub>33</sub>-Trp-Leu-Xaa<sub>37</sub>-Lys-Gly-Arg-Xaa<sub>37</sub>

Formula IX (SEQ ID NO: 12)

- 29 -

wherein:

- Xaa<sub>7</sub> is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine,  
5 homohistidine,  $\alpha$ -fluoromethyl-histidine, or  $\alpha$ -methyl-histidine;  
Xaa<sub>9</sub> is: Ala, Gly, Val, Leu, Ile, Ser, or Thr;  
Xaa<sub>12</sub> is: Phe, Trp, or Tyr;  
Xaa<sub>16</sub> is: Val, Trp, Ile, Leu, Phe, or Tyr;  
10 Xaa<sub>18</sub> is: Ser, Trp, Tyr, Phe, Lys, Ile, Leu, Val;  
Xaa<sub>19</sub> is: Tyr, Trp, or Phe;  
Xaa<sub>20</sub> is: Leu, Phe, Tyr, or Trp;  
Xaa<sub>22</sub> is: Gly, Glu, Asp, or Lys;  
Xaa<sub>25</sub> is: Ala, Val, Ile, or Leu;  
15 Xaa<sub>27</sub> is: Glu, Ile, or Ala;  
Xaa<sub>30</sub> is: Ala or Glu  
Xaa<sub>33</sub> is: Val, or Ile; and  
Xaa<sub>37</sub> is: Gly, His, NH<sub>2</sub>, or is absent.
- 20 Some preferred GLP-1 compounds of formula IX include  
GLP-1(7-37)OH, GLP-1(7-36)-NH<sub>2</sub>, Gly<sup>8</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-GLP-1(7-36)NH<sub>2</sub>, Leu<sup>8</sup>-GLP-1(7-37)OH, Leu<sup>8</sup>-GLP-1(7-36)NH<sub>2</sub>, Ile<sup>8</sup>-GLP-1(7-37)OH, Ile<sup>8</sup>-GLP-1(7-36)NH<sub>2</sub>, Ser<sup>8</sup>-GLP-1(7-37)OH, Ser<sup>8</sup>-GLP-1(7-36)NH<sub>2</sub>,  
25 Thr<sup>8</sup>-GLP-1(7-37)OH, Thr<sup>8</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-Tyr<sup>12</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Tyr<sup>12</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-Tyr<sup>16</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Tyr<sup>16</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Gly<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Gly<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Leu<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Leu<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Ile<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-

- 30 -

37)OH, Ile<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Leu<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-37)OH,  
 Leu<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Ile<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-37)OH, Ile<sup>8</sup>-  
 Asp<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Leu<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-37)OH, Leu<sup>8</sup>-Lys<sup>22</sup>-GLP-  
 1(7-36)NH<sub>2</sub>, Ile<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-37)OH, Ile<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-  
 5 36)NH<sub>2</sub>, Ser<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Ser<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>,  
 Thr<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Thr<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Ser<sup>8</sup>-  
 Asp<sup>22</sup>-GLP-1(7-37)OH, Ser<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Thr<sup>8</sup>-Asp<sup>22</sup>-GLP-  
 1(7-37)OH, Thr<sup>8</sup>-Asp<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Ser<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-  
 37)OH, Ser<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Thr<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-37)OH,  
 10 Thr<sup>8</sup>-Lys<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Glu<sup>22</sup>-GLP-1(7-37)OH, Glu<sup>22</sup>-GLP-1(7-  
 36)NH<sub>2</sub>, Asp<sup>22</sup>-GLP-1(7-37)OH, Asp<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Lys<sup>22</sup>-GLP-  
 1(7-37)OH, Lys<sup>22</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-Ala<sup>27</sup>-GLP-1(7-37)OH,  
 Val<sup>8</sup>-Glu<sup>22</sup>-Ala<sup>27</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-  
 Glu<sup>30</sup>-GLP-1(7-36)NH<sub>2</sub>, Gly<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-37)OH, Gly<sup>8</sup>-Glu<sup>30</sup>-GLP-  
 15 1(7-36)NH<sub>2</sub>, Leu<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-37)OH, Leu<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-  
 36)NH<sub>2</sub>, Ile<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-37)OH, Ile<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-36)NH<sub>2</sub>,  
 Ser<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-37)OH, Ser<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-36)NH<sub>2</sub>, Thr<sup>8</sup>-  
 Glu<sup>30</sup>-GLP-1(7-37)OH, Thr<sup>8</sup>-Glu<sup>30</sup>-GLP-1(7-36)NH<sub>2</sub>, Val<sup>8</sup>-His<sup>37</sup>-GLP-  
 1(7-37)OH, Val<sup>8</sup>-His<sup>37</sup>-GLP-1(7-36)NH<sub>2</sub>, Gly<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH,  
 20 Gly<sup>8</sup>-His<sup>37</sup>-GLP-1(7-36)NH<sub>2</sub>, Leu<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH, Leu<sup>8</sup>-His<sup>37</sup>-  
 GLP-1(7-36)NH<sub>2</sub>, Ile<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH, Ile<sup>8</sup>-His<sup>37</sup>-GLP-1(7-  
 36)NH<sub>2</sub>, Ser<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH, Ser<sup>8</sup>-His<sup>37</sup>-GLP-1(7-36)NH<sub>2</sub>,  
 Thr<sup>8</sup>-His<sup>37</sup>-GLP-1(7-37)OH, Thr<sup>8</sup>-His<sup>37</sup>-GLP-1(7-36)NH<sub>2</sub>.

Some preferred GLP-1 compounds of formula IX having  
 25 multiple substitutions include GLP-1(7-37)OH wherein  
 position 8 is valine or glycine, position 22 is glutamic  
 acid, position 16 is tyrosine, leucine or tryptophan,  
 position 18 is tyrosine, tryptophan, or isoleucine, position  
 25 is valine and position 33 is isoleucine. Other preferred  
 30 GLP-1 compounds include the following: Val<sup>8</sup>-Tyr<sup>16</sup>-GLP-1(7-  
 37)OH, Val<sup>8</sup>-Tyr<sup>12</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Tyr<sup>16</sup>-Phe<sup>19</sup>-GLP-  
 1(7-37)OH, Val<sup>8</sup>-Tyr<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Trp<sup>16</sup>-Glu<sup>22</sup>-  
 GLP-1(7-37)OH, Val<sup>8</sup>-Leu<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Ile<sup>16</sup>-

- 31 -

Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Phe<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH; Val<sup>8</sup>-Trp<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Tyr<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, Val<sup>8</sup>-Phe<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH, and Val<sup>8</sup>-Ile<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37)OH.

- 5           The GLP-1 compounds of the present invention also encompass Exendin compounds. Exendin-3 and Exendin-4 are biologically active peptides first isolated from Helodermatidae lizard venoms and have been shown to bind the GLP-1 receptor and stimulate cAMP-dependent H<sup>+</sup> production in  
10   mammalian parietal cells. Exendin-3 and Exendin-4 are both 39 amino acid peptides which are approximately 53% homologous to GLP-1. They act as potent agonists of GLP-1 activity. Notably, an N-terminally truncated derivative of Exendin, known as Exendin(9-39 amino acids), is an inhibitor  
15   of Exendin-3, Exendin-4 and GLP-1.

          An Exendin compound typically comprises a polypeptide having the amino acid sequence of Exendin-3, Exendin-4, or an analog or fragment thereof. Exendin-3 and Exendin-4 are disclosed in U.S. Patent No. 5,424,286.  
20

          Exendin-3 has the amino acid sequence of SEQ ID NO: 9:

	7	8	9	10	11	12	13	14	15	16	17	
	His	Ser	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	
	18	19	20	21	22	23	24	25	26	27	28	
25	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	
	29	30	31	32	33	34	35	36	37	38	39	
	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	
	40	41	42	43	44	45						
	Gly	Ala	Pro	Pro	Pro	Ser						

30           (SEQ ID NO: 9)

- 32 -

Exendin-4 has the amino acid sequence of SEQ ID NO: 10:

	7	8	9	10	11	12	13	14	15	16	17	
	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	
5	18	19	20	21	22	23	24	25	26	27	28	
	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	
	29	30	31	32	33	34	35	36	37	38	39	
	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	
	40	41	42	43	44	45						
10		Gly	Ala	Pro	Pro	Pro	Ser					
	(SEQ ID NO: 10)											

GLP-1 compounds also include Exendin fragments which are polypeptides obtained after truncation of one or more amino acids from the *N*-terminus and/or C-terminus of Exendin or an Exendin analog. Furthermore, GLP-1 compounds include Exendin polypeptides in which one or more amino acids have been added to the *N*-terminus and/or C-terminus of Exendin or fragments thereof. Exendin compounds of this type have up to about forty-five amino acids.

GLP-1 compounds also include "Exendin analogs." An Exendin analog has sufficient homology to Exendin-4, Exendin-3, or a fragment thereof such that the analog has insulinotropic activity. The activity of Exendin fragments and/or analogs can be assessed using *in vitro* assays such as those described in EP 619,322 and U.S. Patent No. 5,120,712.

Preferably, an Exendin analog has the amino acid sequence of Exendin-4 or a fragment thereof, modified so that from one, two, three, four or five amino acids differ from the amino acid in corresponding position of Exendin-4 or the fragment of Exendin-4. In the nomenclature used herein to designate Exendin compounds, the substituting amino acid and its position is indicated prior to the parent

- 33 -

structure. For example, Val<sup>8</sup>-Exendin-4 designates an Exendin compound in which the glycine normally found at position 8 of Exendin-4 has been replaced with valine.

Another preferred group of GLP-1 compounds is composed of GLP-1/Exendin-4 analogs of formula VIII (SEQ ID NO:11).

```

      7  8  9  10 11 12 13 14 15 16 17
Xaa-Xaa-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Xaa-Ser-
10 18 19 20 21 22 23 24 25 26 27 28
Xaa-Xaa-Xaa-Glu-Xaa-Xaa-Ala-Xaa-Xaa-Xaa-Phe-
29 30 31 32 33 34 35 36 37
Ile-Xaa-Trp-Leu-Xaa-Xaa-Gly-Xaa-R
      formula VIII (SEQ ID NO: 11)
```

15 wherein:

Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;

20 Xaa at position 8 is: Gly, Ala, or Val;  
Xaa at position 16 is: Leu or Val;  
Xaa at position 18 is Lys or Ser;  
Xaa at position 19 is: Gln or Tyr;  
Xaa at position 20 is: Met or Leu;

25 Xaa at position 22 is: Glu or Gln;  
Xaa at position 23 is: Glu, or Gln;  
Xaa at position 25 is: Val or Ala;  
Xaa at position 26 is: Arg or Lys;  
Xaa at position 27 is Leu or Glu;

30 Xaa at position 30 is: Glu or Ala;  
Xaa at position 33 is: Val or Lys;  
Xaa at position 34 is: Asn or Lys;  
Xaa at position 36 is: Gly or Arg; and

- 34 -

R at position 37 is: Gly, Pro, Pro-Ser-Ser-Gly-Ala-Pro-Pro-Ser, or is absent. The activity of 18 different species that fall within this genus is provided in Table 6.

Further Exendin-analogs that are useful for the present invention are described in PCT patent publications WO 99/25728 (Beeley et al.), WO 99/25727 (Beeley et al.), WO 98/05351 (Young et al.), WO 99/40788 (Young et al.), WO 99/07404 (Beeley et al), and WO 99/43708 (Knudsen et al).

The GLP-1 fusion proteins of the present invention can comprise glycosylation sites. Glycosylation is a chemical modification wherein sugar moieties are added to the protein at specific sites. Glycosylation of proteins play a role in ensuring the correct charge, conformation, and stability of maturing protein and can target the protein to the cell surface and eventual secretion of the protein. Most importantly, glycosylation effects the in vivo clearance rate for many proteins. Sugars can be O-linked or N-linked. Generally, O-linked sugars are added to the hydroxyl-group oxygen of serine and threonine, while N-linked sugars are added to the amide nitrogen of asparagine. The consensus site for N-glycosylation is Asn X1 X2 wherein X1 is any amino acid except Pro and X2 is Ser or Thr.

GLP-1 compounds are generally not glycosylated in vivo; however, interestingly the GLP-1 fusion proteins of the present invention that comprise a GLP-1 compound with a C terminal extension fused to an Fc sequence is glycosylated at the last serine in the C terminal extension (SSGAPPPS\*) and at threonine at position 11 in the N terminal region of Fc (AEPKSCDKTHT\*CPFC . . .).

30

- 35 -

Heterologous Fc fusion proteins:

The GLP-1 compounds described above can be fused directly or via a peptide linker to the Fc portion of an immunoglobulin.

Immunoglobulins are molecules containing polypeptide chains held together by disulfide bonds, typically having two light chains and two heavy chains. In each chain, one domain (V) has a variable amino acid sequence depending on the antibody specificity of the molecule. The other domains (C) have a rather constant sequence common to molecules of the same class.

As used herein, the Fc portion of an immunoglobulin has the meaning commonly given to the term in the field of immunology. Specifically, this term refers to an antibody fragment which is obtained by removing the two antigen binding regions (the Fab fragments) from the antibody. One way to remove the Fab fragments is to digest the immunoglobulin with papain protease. Thus, the Fc portion is formed from approximately equal sized fragments of the constant region from both heavy chains, which associate through non-covalent interactions and disulfide bonds. The Fc portion can include the hinge regions and extend through the CH2 and CH3 domains to the C-terminus of the antibody. Representative hinge regions for human and mouse immunoglobulins can be found in Antibody Engineering, A Practical Guide, Borrebaeck, C.A.K., ed., W.H. Freeman and Co., 1992, the teachings of which are herein incorporated by reference. The Fc portion can further include one or more glycosylation sites. The amino acid sequence of a representative Fc protein containing a hinge region, CH2 and CH3 domains, and one N-glycosylation site at position 82 is shown in Figure 1.



- 36 -

There are five types of human immunoglobulin Fc regions with different effect or and pharmacokinetic properties: IgG, IgA, IgM, IgD, and IgE. IgG is the most abundant immunoglobulin in serum. IgG also has the longest half-life  
5 in serum of any immunoglobulin (23 days). Unlike other immunoglobulins, IgG is efficiently recirculated following binding to an Fc receptor. There are four IgG subclasses G1, G2, G3, and G4, each of which have different effect or functions. G1, G2, and G3 can bind C1q and fix complement  
10 while G4 cannot. Even though G3 is able to bind C1q more efficiently than G1, G1 is more effective at mediating complement-directed cell lysis. G2 fixes complement very inefficiently. The C1q binding site in IgG is located at the carboxy terminal region of the CH2 domain.

15 All IgG subclasses are capable of binding to Fc receptors (CD16, CD32, CD64) with G1 and G3 being more effective than G2 and G4. The Fc receptor binding region of IgG is formed by residues located in both the hinge and the carboxy terminal regions of the CH2 domain.

20 IgA can exist both in a monomeric and dimeric form held together by a J-chain. IgA is the second most abundant Ig in serum, but it has a half-life of only 6 days. IgA has three effect or functions. It binds to an IgA specific receptor on macrophages and eosinophils, which drives  
25 phagocytosis and degranulation, respectively. It can also fix complement via an unknown alternative pathway.

IgM is expressed as either a pentamer or a hexamer, both of which are held together by a J-chain. IgM has a serum half-life of 5 days. It binds weakly to C1q via a  
30 binding site located in its CH3 domain. IgD has a half-life of 3 days in serum. It is unclear what effect or functions are attributable to this Ig. IgE is a monomeric Ig and has a serum half-life of 2.5 days. IgE binds to two Fc receptors which drives degranulation and results in the  
35 release of proinflammatory agents.

- 37 -

Depending on the desired *in vivo* effect, the heterologous fusion proteins of the present invention may contain any of the isotypes described above or may contain mutated Fc regions wherein the complement and/or Fc receptor binding functions have been altered. Thus, the heterologous fusion proteins of the present invention may contain the entire Fc portion of an immunoglobulin, fragments of the Fc portion of an immunoglobulin, or analogs thereof fused to a GLP-1 compound.

The fusion proteins of the present invention can consist of single chain proteins or as multi-chain polypeptides. Two or more Fc fusion proteins can be produced such that they interact through disulfide bonds that naturally form between Fc regions. These multimers can be homogeneous with respect to the GLP-1 compound or they may contain different GLP-1 compounds fused at the N-terminus of the Fc portion of the fusion protein.

Regardless of the final structure of the fusion protein, the Fc or Fc-like region must serve to prolong the *in vivo* plasma half-life of the GLP-1 compound fused at the N-terminus. Furthermore, the fused GLP-1 compound must retain some biological activity. An increase in half-life can be demonstrated using the method described in Example 7 wherein the half-life of the fusion protein is compared to the half-life of the GLP-1 compound alone. Biological activity can be determined by *in vitro* and *in vivo* methods known in the art. Representative biological assays are described in Examples 6, 8, and 9.

Since the Fc region of IgG produced by proteolysis has the same *in vivo* half-life as the intact IgG molecule and Fab fragments are rapidly degraded, it is believed that the relevant sequence for prolonging half-life reside in the CH2 and/or CH3 domains. Further, it has been shown in the literature that the catabolic rates of IgG variants that do not bind the high-affinity Fc receptor or C1q are

- 38 -

indistinguishable from the rate of clearance of the parent wild-type antibody, indicating that the catabolic site is distinct from the sites involved in Fc receptor or C1q binding. [Nawrzynczak et al., (1992) *Molecular Immunology* 29:221]. Site-directed mutagenesis studies using a murine IgG1 Fc region suggested that the site of the IgG1 Fc region that controls the catabolic rate is located at the CH2-CH3 domain interface.

Based on these studies, Fc regions can be modified at the catabolic site to optimize the half-life of the fusion proteins. It is preferable that the Fc region used for the heterologous fusion proteins of the present invention be derived from an IgG1 or an IgG4 Fc region. It is even more preferable that the Fc region be IgG4 or derived from IgG4. Preferably the IgG Fc region contains both the CH2 and CH3 regions including the hinge region.

#### Heterologous albumin fusion proteins:

The GLP-1 compounds described above can be fused directly or via a peptide linker to albumin or an analog, fragment, or derivative thereof.

Generally the albumin proteins making up part of the fusion proteins of the present invention can be derived from albumin cloned from any species. However, human albumin and fragments and analogs thereof are preferred to reduce the risk of the fusion protein being immunogenic in humans. Human serum albumin (HSA) consists of a single non-glycosylated polypeptide chain of 585 amino acids with a formula molecular weight of 66,500. The amino acid sequence of human HSA is shown in figure 2. [See Meloun, et al. (1975) *FEBS Letters* 58:136; Behrens, et al. (1975) *Fed. Proc.* 34:591; Lawn, et al. (1981) *Nucleic Acids Research* 9:6102-6114; Minghetti, et al. (1986) *J. Biol. Chem.* 261:6747]. A variety of polymorphic variants as well as analogs and fragments of albumin have been described. [See

- 39 -

Weitkamp, et al., (1973) Ann. Hum. Genet. 37:219]. For example, in EP 322,094, the inventors disclose various shorter forms of HSA. Some of these fragments include HSA(1-373), HSA(1-388), HSA(1-389), HSA(1-369), and HSA(1-419) and fragments between 1-369 and 1-419. EP 399,666 discloses albumin fragments that include HSA(1-177) and HSA(1-200) and fragments between HSA(1-177) and HSA(1-200).

It is understood that the heterologous fusion proteins of the present invention include GLP-1 compounds that are coupled to any albumin protein including fragments, analogs, and derivatives wherein such fusion protein is biologically active and has a longer plasma half-life than the GLP-1 compound alone. Thus, the albumin portion of the fusion protein need not necessarily have a plasma half-life equal to that of native human albumin. Fragments, analogs, and derivatives are known or can be generated that have longer half-lives or have half-lives intermediate to that of native human albumin and the GLP-1 compound of interest.

The heterologous fusion proteins of the present invention encompass proteins having conservative amino acid substitutions in the GLP-1 compound and/or the Fc or albumin portion of the fusion protein. A "conservative substitution" is the replacement of an amino acid with another amino acid that has the same net electronic charge and approximately the same size and shape. Amino acids with aliphatic or substituted aliphatic amino acid side chains have approximately the same size when the total number carbon and heteroatoms in their side chains differs by no more than about four. They have approximately the same shape when the number of branches in their side chains differs by no more than one. Amino acids with phenyl or substituted phenyl groups in their side chains are considered to have about the same size and shape. Except as otherwise specifically provided herein, conservative

- 40 -

substitutions are preferably made with naturally occurring amino acids.

However, the term "amino acid" is used herein in its broadest sense, and includes naturally occurring amino acids as well as non-naturally occurring amino acids, including amino acid analogs and derivatives. The latter includes molecules containing an amino acid moiety. One skilled in the art will recognize, in view of this broad definition, that reference herein to an amino acid includes, for example, naturally occurring proteogenic L-amino acids; D-amino acids; chemically modified amino acids such as amino acid analogs and derivatives; naturally occurring non-proteogenic amino acids such as norleucine,  $\beta$ -alanine, ornithine, GABA, etc.; and chemically synthesized compounds having properties known in the art to be characteristic of amino acids. As used herein, the term "proteogenic" indicates that the amino acid can be incorporated into a peptide, polypeptide, or protein in a cell through a metabolic pathway.

The incorporation of non-natural amino acids, including synthetic non-native amino acids, substituted amino acids, or one or more D-amino acids into the heterologous fusion proteins of the present invention can be advantageous in a number of different ways. D-amino acid-containing peptides, etc., exhibit increased stability *in vitro* or *in vivo* compared to L-amino acid-containing counterparts. Thus, the construction of peptides, etc., incorporating D-amino acids can be particularly useful when greater intracellular stability is desired or required. More specifically, D-peptides, etc., are resistant to endogenous peptidases and proteases, thereby providing improved bioavailability of the molecule, and prolonged lifetimes *in vivo* when such properties are desirable. Additionally, D-peptides, etc., cannot be processed efficiently for major histocompatibility complex class II-restricted presentation to T helper cells,

- 41 -

and are therefore, less likely to induce humoral immune responses in the whole organism.

In addition to structure/function analyses of the various polypeptides encompassed by the present invention, there are numerous factors that can be considered when selecting amino acids for substitution. One factor that can be considered in making such changes is the hydropathic index of amino acids. The importance of the hydropathic amino acid index in conferring interactive biological function on a protein has been discussed by Kyte and Doolittle (1982, *J. Mol. Biol.*, 157: 105-132). It is accepted that the relative hydropathic character of amino acids contributes to the secondary structure of the resultant protein. This, in turn, affects the interaction of the protein with molecules such as enzymes, substrates, receptors, ligands, DNA, antibodies, antigens, etc. Based on its hydrophobicity and charge characteristics, each amino acid has been assigned a hydropathic index as follows:

isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate/glutamine/aspartate/asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

As is known in the art, certain amino acids in a peptide, polypeptide, or protein can be substituted for other amino acids having a similar hydropathic index or score and produce a resultant peptide, etc., having similar or even improved biological activity. In making such changes, it is preferable that amino acids having hydropathic indices within  $\pm 2$  are substituted for one another. More preferred substitutions are those wherein the amino acids have hydropathic indices within  $\pm 1$ . Most

- 42 -

preferred substitutions are those wherein the amino acids have hydropathic indices within  $\pm 0.5$ .

Like amino acids can also be substituted on the basis of hydrophilicity. U.S. Patent No. 4,554,101 discloses that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein. The following hydrophilicity values have been assigned to amino acids: arginine/lysine (+3.0); aspartate/glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine/glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 $\pm$ 1); alanine/histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine/isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); and tryptophan (-3.4). Thus, one amino acid in a peptide, polypeptide, or protein can be substituted by another amino acid having a similar hydrophilicity score and still produce a resultant peptide, etc., having similar biological activity, i.e., still retaining correct biological function. In making such changes, amino acids having hydropathic indices within  $\pm 2$  are preferably substituted for one another, those within  $\pm 1$  are more preferred, and those within  $\pm 0.5$  are most preferred.

As outlined above, amino acid substitutions in the fusion proteins of the present invention can be based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, etc. Furthermore, substitutions can be made based on secondary structure propensity. For example, a helical amino acid can be replaced with an amino acid that would preserve the helical structure. Exemplary substitutions that take various of the foregoing characteristics into consideration in order to produce conservative amino acid changes resulting in silent changes within the present peptides, etc., can be selected from other members of the class to which the naturally

- 43 -

occurring amino acid belongs. Amino acids can be divided into the following four groups: (1) acidic amino acids; (2) basic amino acids; (3) neutral polar amino acids; and (4) neutral non-polar amino acids.

5

General methods for making the heterologous fusion proteins of the present invention.

Although the heterologous fusion proteins of the present invention can be made by a variety of different methods, recombinant methods are preferred. For purposes of the present invention, as disclosed and claimed herein, the following general molecular biology terms and abbreviations are defined below. The terms and abbreviations used in this document have their normal meanings unless otherwise designated. For example, "°C" refers to degrees Celsius; "mmol" refers to millimole or millimoles; "mg" refers to milligrams; "µg" refers to micrograms; "ml or mL" refers to milliliters; and "µl or µL" refers to microliters. Amino acids abbreviations are as set forth in 37 C.F.R. § 1.822 (b)(2) (1994).

"Base pair" or "bp" as used herein refers to DNA or RNA. The abbreviations A,C,G, and T correspond to the 5'-monophosphate forms of the deoxyribonucleosides (deoxy)adenosine, (deoxy)cytidine, (deoxy)guanosine, and thymidine, respectively, when they occur in DNA molecules. The abbreviations U,C,G, and A correspond to the 5'-monophosphate forms of the ribonucleosides uridine, cytidine, guanosine, and adenosine, respectively when they occur in RNA molecules. In double stranded DNA, base pair may refer to a partnership of A with T or C with G. In a DNA/RNA, heteroduplex base pair may refer to a partnership of A with U or C with G. (See the definition of "complementary", *infra*.)

"Digestion" or "Restriction" of DNA refers to the catalytic cleavage of the DNA with a restriction enzyme that



- 44 -

acts only at certain sequences in the DNA ("sequence-specific endonucleases"). The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors, and other requirements were used as would be known to one of ordinary skill in the art. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer or can be readily found in the literature.

10 "Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments. Unless otherwise provided, ligation may be accomplished using known buffers and conditions with a DNA ligase, such as T4 DNA ligase.

15 "Plasmid" refers to an extrachromosomal (usually) self-replicating genetic element. Plasmids are generally designated by a lower case "p" followed by letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available 20 plasmids in accordance with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

25 "Recombinant DNA cloning vector" as used herein refers to any autonomously replicating agent, including, but not limited to, plasmids and phages, comprising a DNA molecule to which one or more additional DNA segments can or have been added.

30 "Recombinant DNA expression vector" as used herein refers to any recombinant DNA cloning vector in which a promoter to control transcription of the inserted DNA has been incorporated.

"Transcription" refers to the process whereby information contained in a nucleotide sequence of DNA is 35 transferred to a complementary RNA sequence.

- 45 -

"Transfection" refers to the uptake of an expression vector by a host cell whether or not any coding sequences are, in fact, expressed. Numerous methods of transfection are known to the ordinarily skilled artisan, for example, calcium phosphate co-precipitation, liposome transfection, and electroporation. Successful transfection is generally recognized when any indication of the operation of this vector occurs within the host cell.

"Transformation" refers to the introduction of DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integration. Methods of transforming bacterial and eukaryotic hosts are well known in the art, many of which methods, such as nuclear injection, protoplast fusion or by calcium treatment using calcium chloride are summarized in J. Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, (1989). Generally, when introducing DNA into Yeast the term transformation is used as opposed to the term transfection.

"Translation" as used herein refers to the process whereby the genetic information of messenger RNA (mRNA) is used to specify and direct the synthesis of a polypeptide chain.

"Vector" refers to a nucleic acid compound used for the transfection and/or transformation of cells in gene manipulation bearing polynucleotide sequences corresponding to appropriate protein molecules which, when combined with appropriate control sequences, confers specific properties on the host cell to be transfected and/or transformed. Plasmids, viruses, and bacteriophage are suitable vectors. Artificial vectors are constructed by cutting and joining DNA molecules from different sources using restriction enzymes and ligases. The term "vector" as used herein includes Recombinant DNA cloning vectors and Recombinant DNA expression vectors.

- 46 -

"Complementary" or "Complementarity", as used herein, refers to pairs of bases (purines and pyrimidines) that associate through hydrogen bonding in a double stranded nucleic acid. The following base pairs are complementary:  
5 guanine and cytosine; adenine and thymine; and adenine and uracil.

"Hybridization" as used herein refers to a process in which a strand of nucleic acid joins with a complementary strand through base pairing. The conditions employed in the  
10 hybridization of two non-identical, but very similar, complementary nucleic acids varies with the degree of complementarity of the two strands and the length of the strands. Such techniques and conditions are well known to practitioners in this field.

15 "Isolated amino acid sequence" refers to any amino acid sequence, however, constructed or synthesized, which is locationally distinct from the naturally occurring sequence.

"Isolated DNA compound" refers to any DNA sequence, however constructed or synthesized, which is locationally  
20 distinct from its natural location in genomic DNA.

"Isolated nucleic acid compound" refers to any RNA or DNA sequence, however constructed or synthesized, which is locationally distinct from its natural location.

25 "Primer" refers to a nucleic acid fragment which functions as an initiating substrate for enzymatic or synthetic elongation.

"Promoter" refers to a DNA sequence which directs transcription of DNA to RNA.

30 "Probe" refers to a nucleic acid compound or a fragment, thereof, which hybridizes with another nucleic acid compound.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe  
35 length, washing temperature, and salt concentration. In

- 47 -

general, longer probes require higher temperatures for proper annealing, while short probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to re-anneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature that can be used. As a result, it follows that higher relative temperatures would tend to make the reactions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, 1995.

"Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that (1) employ low ionic strength and high temperature for washing, for example, 15 mM sodium chloride/1.5 mM sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride/75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5X SSC (750 mM sodium chloride, 75 mM sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5X Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C with washes at 42°C in 0.2X SSC (30 mM sodium chloride/3 mM sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1X SSC containing EDTA at 55°C.

"Moderately stringent conditions" may be identified as described by Sambrook et al. [*Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, (1989)], and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength, and %SDS) less

- 48 -

stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5X SSC (750 mM sodium chloride, 75 mM sodium citrate), 50 mM sodium phosphate at pH 7.6, 5X Denhardt's solution, 10% dextran sulfate, and 20 mg/mL denatured sheared salmon sperm DNA, followed by washing the filters in 1X SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc., as necessary to accommodate factors such as probe length and the like.

"PCR" refers to the widely-known polymerase chain reaction employing a thermally-stable DNA polymerase.

"Leader sequence" refers to a sequence of amino acids which can be enzymatically or chemically removed to produce the desired polypeptide of interest.

"Secretion signal sequence" refers to a sequence of amino acids generally present at the N-terminal region of a larger polypeptide functioning to initiate association of that polypeptide with the cell membrane compartments like endoplasmic reticulum and secretion of that polypeptide through the plasma membrane.

Construction of DNA encoding the heterologous fusion proteins of the present invention:

Wild-type albumin and Immunoglobulin proteins can be obtained from a variety of sources. For example, these proteins can be obtained from a cDNA library prepared from tissue or cells which express the mRNA of interest at a detectable level. Libraries can be screened with probes designed using the published DNA or protein sequence for the particular protein of interest. For example, immunoglobulin light or heavy chain constant regions are described in Adams, et al. (1980) Biochemistry 19:2711-2719; Goughet, et al. (1980) Biochemistry 19:2702-2710; Dolby, et al. (1980) Proc. Natl. Acad. Sci. USA 77:6027-6031; Rice et al. (1982)

- 49 -

Proc. Natl. Acad. Sci. USA 79:7862-7862; Palkner, et al. (1982) Nature 298:286-288; and Morrison, et al. (1984) Ann. Rev. Immunol. 2:239-256. Some references disclosing albumin protein and DNA sequences include Meloun, et al. (1975) FEBS Letters 58:136; Behrens, et al. (1975) Fed. Proc. 34:591; Lawn, et al. (1981) Nucleic Acids Research 9:6102-6114; and Minghetti, et al. (1986) J. Biol. Chem. 261:6747

Screening a cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1989). An alternative means to isolate a gene encoding an albumin or immunoglobulin protein is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1995)]. PCR primers can be designed based on published sequences.

Generally the full-length wild-type sequences cloned from a particular species can serve as a template to create analogs, fragments, and derivatives that retain the ability to confer a longer plasma half-life on the GLP-1 compound that is part of the fusion protein. It is preferred that the Fc and albumin portions of the heterologous fusion proteins of the present invention be derived from the native human sequence in order to reduce the risk of potential immunogenicity of the fusion protein in humans.

In particular, it is preferred that the immunoglobulin portion of a fusion protein encompassed by the present invention contain only an Fc fragment of the immunoglobulin.

Depending on whether particular effect or functions are desired and the structural characteristics of the fusion protein, an Fc fragment may contain the hinge region along with the CH2 and CH3 domains or some other combination thereof. These Fc fragments can be generated using PCR techniques with primers designed to hybridize to sequences

- 50 -

corresponding to the desired ends of the fragment. Similarly, if fragments of albumin are desired, PCR primers can be designed which are complementary to internal albumin sequences. PCR primers can also be designed to create  
5 restriction enzyme sites to facilitate cloning into expression vectors.

DNA encoding the GLP-1 compounds of the present invention can be made by a variety of different methods including cloning methods like those described above as well  
10 as chemically synthesized DNA. Chemical synthesis may be attractive given the short length of the encoded peptide. The amino acid sequence for GLP-1 has been published as well as the sequence of the preproglucagon gene. [Lopez, et al. (1983) Proc. Natl. Acad. Sci., USA 80:5485-5489; Bell, et  
15 al. (1983) Nature, 302:716-718; Heinrich, G., et al. (1984) Endocrinol, 115:2176-2181; Ghiglione, M., et al. 91984) Diabetologia 27:599-600]. Thus, primers can be designed to PCR native GLP-1 compounds and fragments thereof.

The gene encoding a fusion protein can then be  
20 constructed by ligating DNA encoding a GLP-1 compound in-frame to DNA encoding an albumin or Fc protein. The gene encoding the GLP-1 compound and the gene encoding the albumin or Fc protein can also be joined in-frame via DNA encoding a linker peptide.

The *in vivo* function and stability of the heterologous fusion proteins of the present invention can be optimized by adding small peptide linkers to prevent potentially unwanted domain interactions. Although these linkers can potentially be any length and consist of any combination of amino acids,  
25 it is preferred that the length be no longer than necessary to prevent unwanted domain interactions and/or optimize biological activity and/or stability. Generally, the linkers should not contain amino acids with extremely bulky side chains or amino acids likely to introduce significant  
30 secondary structure. It is preferred that the linker be  
35

- 51 -

serine-glycine rich and be less than 30 amino acids in length. It is more preferred that the linker be no more than 20 amino acids in length. It is even more preferred that the linker be no more than 15 amino acids in length. A preferred linker contains repeats of the sequence Gly-Gly-Gly-Gly-Ser. It is preferred that there be between 2 and 6 repeats of this sequence. It is even more preferred that there be between 3 and 4 repeats of this sequence.

The DNA encoding wild-type GLP-1, albumin, and Fc polypeptides and fragments thereof can be mutated either before ligation or in the context of a cDNA encoding an entire fusion protein. A variety of mutagenesis techniques are well known in the art. For example, a mutagenic PCR method utilizes strand overlap extension to create specific base mutations for the purposes of changing a specific amino acid sequence in the corresponding protein. This PCR mutagenesis requires the use of four primers, two in the forward orientation (primers A and C) and two in the reverse orientation (primers B and D). A mutated gene is amplified from the wild-type template in two different stages. The first reaction amplifies the gene in halves by performing an A to B reaction and a separate C to D reaction wherein the B and C primers target the area of the gene to be mutated. When aligning these primers with the target area, they contain mismatches for the bases that are targeted to be changed. Once the A to B and C to D reactions are complete, the reaction products are isolated and mixed for use as the template for the A to D reaction. This reaction then yields the full, mutated product.

Once a gene encoding an entire fusion protein is produced it can be cloned into an appropriate expression vector. Specific strategies that can be employed to make the GLP-1 fusion proteins of the present invention are described in example 1.

35



- 52 -

General methods to recombinantly express the heterologous fusion proteins of the present invention:

Host cells are transfected or transformed with expression or cloning vectors described herein for heterologous fusion protein production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: A Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook, et al., supra. Methods of transfection are known to the ordinarily skilled artisan, for example,  $\text{CaPO}_4$  and electroporation. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of van Solingen et al., *J Bact.* 130(2): 946-7 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. USA* 76(8): 3829-33 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene or polyomithine, may also be used. For various techniques for transforming mammalian cells, see Keown, et al., *Methods in Enzymology* 185: 527-37 (1990) and Mansour, et al., *Nature* 336(6197): 348-52 (1988).

Suitable host cells for cloning or expressing the nucleic acid (e.g., DNA) in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 3 1.446); *E. coli* X1 776 (ATCC 3 1.537); *E. coli* strain W3 110

- 53 -

(ATCC 27.325) and K5 772 (ATCC 53.635). Other suitable prokaryotic host cells include Enterobacteriaceae such as Escherichia, e.g., E. coli, Enterobacter, Erwinia, Klebsiella, Proteus, Salmonella, e.g., Salmonella typhimurium, Serratia, e.g., Serratia marcescens, and Shigella, as well as Bacilli such as B. subtilis and B. licheniformis (e.g., B. licheniformis 4 1 P disclosed in DD266,7 10, published 12 April 1989), Pseudomonas such as P. aeruginosa, and Streptomyces. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3 110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including E. coli W3110 strain 1A2, which has the complete genotype ronaA; E. coli W3 110 strain 9E4, which has the complete genotype ton4 ptr3; E. coli W3110 strain 27C7 (ATCC 55,244), which has the complete genotype tonA, ptr3 phoA E15 (argF-lac) I69 degP ompT /can'; E. coli W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant degP deletion mutation; and an E. coli strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, in vivo methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for fusion protein vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* [Beach and Nurse, *Nature* 290: 140-3 (1981); EP 139,383 published 2 May 1995]; *Muyveromyces* hosts [U.S. Patent No. 4,943,529; Fleer, et al., *Bio/Technology* 9(10): 968-75 (1991)] such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574) [de Lourencourt et al., *J.*

- 54 -

- Bacteriol.* 154(2): 737-42 (1983)]; *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906) [Van den Berg et al., *Bio/Technology* 8(2): 135-9 (1990)]; *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070) [Sreekrishna et al., *J. Basic Microbiol.* 28(4): 265-78 (1988)]; *Candida*; *Trichoderma reesei* (EP 244,234); *Neurospora crassa* [Case, et al., *Proc. Natl. Acad. Sci. USA* 76(10): 5259-63 (1979)]; Schwanniomycetes such as
- 10 *Schwanniomycetes occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* [Ballance et al., *Biochem. Biophys. Res. Comm.* 112(1): 284-9 (1983)];
- 15 Tilburn, et al., *Gene* 26(2-3): 205-21 (1983); Yelton, et al., *Proc. Natl. Acad. Sci. USA* 81(5): 1470-4 (1984)] and *A. niger* [Kelly and Hynes, *EMBO J.* 4(2): 475-9 (1985)]. Methylotrophic yeasts are selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and
- 20 *Rhodotorula*. A list of specific species that are exemplary of this class of yeast may be found in C. Antony, *The Biochemistry of Methylotrophs* 269 (1982).

- Suitable host cells for the expression of the fusion proteins of the present invention are derived from
- 25 multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sp, *Spodoptera high5* as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney
- 30 CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line [293 or 293 cells subcloned for growth in suspension culture, Graham, et al., *J. Gen. Virol.*, 36(1): 59-74 (1977)]; Chinese hamster ovary cells/-DHFR [CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77(7): 4216-20
- 35 (1980)]; mouse sertoli cells [TM4, Mather, *Biol. Reprod.*

- 55 -

23(1):243-52 (1980)]; human lung cells (WI38, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

5       The fusion proteins of the present invention may be recombinantly produced directly, or as a protein having a signal sequence or other additional sequences which create a specific cleavage site at the N-terminus of the mature fusion protein. In general, the signal sequence may be a component  
10 of the vector, or it may be a part of the fusion protein-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders.  
15 For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* cc-factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179), or  
20 the signal described in WO 90/13646. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species as well as viral secretory leaders.

25       Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-  
30 negative bacteria, the 2u plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

      Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical  
35 selection genes encode proteins that (a) confer resistance to

- 56 -

antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement autotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the fusion protein-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described [Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77(7): 4216-20 (1980)]. A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb, et al., *Nature* 282(5734): 39-43 (1979); Kingsman, et al., *Gene* 7(2): 141-52 (1979); Tschumper, et al., *Gene* 10(2): 157-66 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP1 [Jones, *Genetics* 85: 23-33 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the fusion protein-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the  $\beta$ -lactamase and lactose promoter systems [Chang, et al., *Nature* 275(5681): 617-24 (1978); Goeddel, et al., *Nature* 281(5732): 544-8 (1979)], alkaline phosphatase, a tryptophan (up) promoter system [Goeddel, *Nucleic Acids Res.* 8(18): 4057-74 (1980); EP 36,776 published 30 September 1981], and hybrid promoters such as the *tat* promoter [deBoer, et al., *Proc. Natl. Acad. Sci. USA* 80(1): 21-5 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the fusion protein.

- 57 -

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman, et al., *J. Biol. Chem.* 255(24): 12073-80 (1980)] or other glycolytic enzymes [Hess et al., *J. Adv. Enzyme Reg.* 7: 149 (1968); Holland, *Biochemistry* 17(23): 4900-7 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657. Transcription of fusion protein-encoding mRNA from vectors in mammalian host cells may be controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a polynucleotide encoding a fusion protein by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin,

- 58 -

elastase, albumin, a-ketoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the fusion protein coding sequence but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and occasionally 3' untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the fusion protein.

Various forms of a fusion protein may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g., Triton-X 100) or by enzymatic cleavage. Cells employed in expression of a fusion protein can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

Purification of the heterologous fusion proteins of the present invention:

Once the heterologous fusion proteins of the present invention are expressed in the appropriate host cell, the analogs can be isolated and purified. The following procedures are exemplary of suitable purification procedures: fractionation on carboxymethyl cellulose; gel filtration such

- 59 -

as Sephadex G-75; anion exchange resin such as DEAE or Mono-Q; cation exchange such as CM or Mono-S; protein A sepharose to remove contaminants such as IgG; metal chelating columns to bind epitope-tagged forms of the polypeptide; reversed-phase HPLC; chromatofocusing; silica gel; ethanol precipitation; and ammonium sulfate precipitation.

Various methods of protein purification may be employed and such methods are known in the art and described, for example, in Deutscher, *Methods in Enzymology* 182: 83-9 (1990) and Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, NY (1982). The purification step(s) selected will depend on the nature of the production process used and the particular fusion protein produced. For example, fusion proteins comprising an Fc fragment can be effectively purified using a Protein A or Protein G affinity matrix. Low or high pH buffers can be used to elute the fusion protein from the affinity matrix. Mild elution conditions will aid in preventing irreversible denaturation of the fusion protein. Imidazole-containing buffers can also be used. Example 3 describes some successful purification protocols for the fusion proteins of the present invention.

Characterization of the heterologous fusion proteins of the present invention:

Numerous methods exist to characterize the fusion proteins of the present invention. Some of these methods include: SDS-PAGE coupled with protein staining methods or immunoblotting using anti-IgG or anti-HSA antibodies. Other methods include matrix assisted laser desorption/ionization-mass spectrometry (MALDI-MS), liquid chromatography/mass spectrometry, isoelectric focusing, analytical anion exchange, chromatofocussing, and circular dichroism to name a few. A representative number of heterologous fusion proteins were characterized using SDS-PAGE coupled with immunoblotting as



- 60 -

well as mass spectrometry (See examples 4 and 5 and figures 3 and 4).

For example table 3 (see example 5) illustrates the calculated molecular mass for a representative number of fusion proteins as well as the mass as determined by mass spectrometry. In addition, Figures 3 and 4 illustrate molecular weights of a representative number of fusion proteins as determined by SDS PAGE. All heterologous fusion proteins tested were expressed and secreted transiently. In addition, the Igk signal sequence was cleaved to yield proteins with the correct N-terminus.

Further, table 3 illustrates that in some instances the mass determined by mass spectrometry is greater than expected. This is the result of glycosylation of the Fc portion and the C terminal extension. Enzymatic digestion of the fusion proteins followed by reversed-phase HPLC and mass spectrometry can identify peptide fractions that contain sugar moieties. These fractions can then be N-terminal amino acid sequenced to identify the potential glycosylation site. For example, characterization of Exendin-4-Fc (SEQ ID NO: 29) shows that the serine at position 39 and threonine at position 50 are O-linked glycosylated and the asparagine at position 122 is N-linked glycosylated.

A representative number of GLP-1 fusion proteins were also tested for activity. Numerous methods exist to detect GLP-1 activity *in vitro* and *in vivo* (see examples 6, 7, 8, and 9). Table 4 (example 6) illustrates GLP-1 receptor activity associated with several GLP-1 fusions. The numbers are relative to the activity associated with Val<sup>6</sup>-GLP-1(7-37)OH. All fusion proteins tested had GLP-1 receptor activity. A low level of *in vitro* activity is not necessarily indicative of a weak effect *in vivo*. Because of the substantial increase in the half-life of these fusion proteins, weak *in vitro* activity is not generally a predictor of weak *in vivo* activity. Figure 7 and example 7 illustrate the prolonged half-life associated

- 61 -

with the fusion proteins of the present invention. For example, Val<sup>8</sup>-GLP-1-Fc had a half-life of approximately 45 hours in monkeys, Val<sup>8</sup>-GLP-1-HSA had a half-life of about 87 hours in monkeys, Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-IgG1 had a half-life after IV administration of approximately 55 hours in dogs, and Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-IgG1 had a half-life after SC administration of approximately 38 hours in dogs.

Compositions of the invention:

Physical stability is also an essential feature for therapeutic protein formulations. GLP-1 compounds have been especially difficult to manufacture and formulate due to structural changes that occur during processing. For example, some GLP-1 compounds have a general tendency to aggregate. In addition, it has been shown that some GLP-1 compounds convert from a soluble and active  $\alpha$ -helix form to an insoluble and potentially inactive  $\beta$ -sheet form. The fusion of GLP-1 compounds to large proteins such as the Fc region of an IgG or albumin not only acts to increase the half-life of the GLP-1 compound, but also contributes to the physical and conformational stability of the GLP-1 compound. For example, Val<sup>8</sup>-GLP-1-Linker-HSA in PBS is stable at 37°C out to about 30 days.

The heterologous fusion proteins of the present invention may be formulated with one or more excipients. The active fusion proteins of the present invention may be combined with a pharmaceutically acceptable buffer, and the pH adjusted to provide acceptable stability, and a pH acceptable for administration such as parenteral administration.

Optionally, one or more pharmaceutically-acceptable antimicrobial agents may be added. Meta-cresol and phenol are preferred pharmaceutically-acceptable microbial agents. One or more pharmaceutically-acceptable salts may be added to adjust the ionic strength or tonicity. One or more excipients may be added to further adjust the isotonicity of the

- 62 -

formulation. Glycerin is an example of an isotonicity-adjusting excipient. Pharmaceutically acceptable means suitable for administration to a human or other animal and thus, does not contain toxic elements or undesirable  
5 contaminants and does not interfere with the activity of the active compounds therein.

A pharmaceutically-acceptable salt form of the heterologous fusion proteins of the present invention may be used in the present invention. Acids commonly employed to  
10 form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydriodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as *p*-toluenesulfonic acid, methanesulfonic acid, oxalic acid, *p*-bromophenyl-sulfonic acid, carbonic acid, succinic acid,  
15 citric acid, benzoic acid, acetic acid, and the like. Preferred acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid.

Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal  
20 hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, and the like.

#### 25 Administration of Compositions:

Administration may be via any route known to be effective by the physician of ordinary skill. Peripheral, parenteral is one such method. Parenteral administration is commonly understood in the medical literature as the injection of a  
30 dosage form into the body by a sterile syringe or some other mechanical device such as an infusion pump. Peripheral parenteral routes can include intravenous, intramuscular, subcutaneous, and intraperitoneal routes of administration.

The heterologous fusion proteins of the present invention  
35 may also be amenable to administration by oral, rectal, nasal,

- 63 -

or lower respiratory routes, which are non-parenteral routes. Of these non-parenteral routes, the lower respiratory route and the oral route are preferred.

The fusion proteins of the present invention can be used  
5 to treat a wide variety of diseases and conditions. The fusion proteins of the present invention primarily exert their biological effects by acting at a receptor referred to as the "GLP-1 receptor." Subjects with diseases and/or conditions that respond favorably to GLP-1 receptor stimulation or to the  
10 administration of GLP-1 compounds can therefore be treated with the GLP-1 fusion proteins of the present invention. These subjects are said to "be in need of treatment with GLP-1 compounds" or "in need of GLP-1 receptor stimulation". Included are subjects with non-insulin dependent diabetes,  
15 insulin dependent diabetes, stroke (see WO 00/16797), myocardial infarction (see WO 98/08531), obesity (see WO 98/19698), catabolic changes after surgery (see U.S. Patent No. 6,006,753), functional dyspepsia and irritable bowel syndrome (see WO 99/64060). Also included are subjects  
20 requiring prophylactic treatment with a GLP-1 compound, e.g., subjects at risk for developing non-insulin dependent diabetes (see WO 00/07617). Subjects with impaired glucose tolerance or impaired fasting glucose, subjects whose body weight is about 25% above normal body weight for the subject's height  
25 and body build, subjects with a partial pancreatectomy, subjects having one or more parents with non-insulin dependent diabetes, subjects who have had gestational diabetes and subjects who have had acute or chronic pancreatitis are at risk for developing non-insulin dependent diabetes.  
30 An "effective amount" of a GLP-1 compound is the quantity which results in a desired therapeutic and/or prophylactic effect without causing unacceptable side-effects when administered to a subject in need of GLP-1 receptor stimulation. A "desired therapeutic effect"  
35 includes one or more of the following: 1) an amelioration of

- 64 -

the symptom(s) associated with the disease or condition; 2) a delay in the onset of symptoms associated with the disease or condition; 3) increased longevity compared with the absence of the treatment; and 4) greater quality of life compared with the absence of the treatment. For example, an "effective amount" of a GLP-1 compound for the treatment of diabetes is the quantity that would result in greater control of blood glucose concentration than in the absence of treatment, thereby resulting in a delay in the onset of diabetic complications such as retinopathy, neuropathy or kidney disease. An "effective amount" of a GLP-1 compound for the prevention of diabetes is the quantity that would delay, compared with the absence of treatment, the onset of elevated blood glucose levels that require treatment with anti-hypoglycaemic drugs such as sulfonyl ureas, thiazolidinediones, insulin and/or biguanidines.

The dose of fusion protein effective to normalize a patient's blood glucose will depend on a number of factors, among which are included, without limitation, the subject's sex, weight and age, the severity of inability to regulate blood glucose, the route of administration and bioavailability, the pharmacokinetic profile of the fusion protein, the potency, and the formulation.

The present invention comprises GLP-1 compounds that have improved biochemical and biophysical properties by virtue of being fused to an albumin protein, an albumin fragment, an albumin analog, a Fc protein, a Fc fragment, or a Fc analog. These heterologous proteins can be successfully expressed in host cells, retain signaling activities associated with activation of the GLP-1 receptor, and have prolonged half-lives.

The following examples are presented to further describe the present invention. The scope of the present invention is not to be construed as merely consisting of the following examples. Those skilled in the art will recognize

- 65 -

that the particular reagents, equipment, and procedures described are merely illustrative and are not intended to limit the present invention in any manner.

5                    Example 1: Construction of DNA  
                  encoding heterologous fusion proteins

Example 1a Construction of DNA encoding Val<sup>8</sup>-GLP-1(7-37)-

        Fc: A Fc portion of human IgG1 was isolated from a  
cDNA library and contains the full hinge region and the CH2  
10 and CH3 domains. A fragment containing 696 base pairs of  
this Fc portion of human IgG1 was subcloned into the *NheI*  
and *Eco47III* sites of mammalian expression vector pJB02 to  
create pJB02/Fc (see Figure 5). DNA encoding the Igk  
secretion signal sequence fused to Val<sup>8</sup>-GLP-1(7-37) was  
15 generated by *in vitro* hybridization of four overlapping and  
complementary oligonucleotides:

5'- CTAGCCACCATGGAGACAGACACACTCCTGCTATGGGTACTGCTGCTCTGGGTT  
CCAGGTTCCACTGGTGACCACTG - 3' [SEQ ID NO:12]

20

5'- GAGGGCACCTTCACCTCCGACGTGTCCTCTATCTGGAGGGCCAGGCCCAAGGA  
GTTTCATCGCCTGGCTGGTGAAGGAAGAGGC - 3' [SEQ ID NO:13]

5'- TGAAGGTGCCTCCACGTGGTCACCACTGGAACCTGGAACCCAGAGCAGCAGTA  
25 CCCATAGCAGGAGTGTGTCGTCTCCATGGTGG - 3' [SEQ ID  
NO:14]

5'- GCCTCTTCCCTTCACCAAGCCAGGCGATGAACTCCTTGGCGGCTGGCCCTCCAGA  
TAGGAGGACACGTCGGAGG - 3' [SEQ ID NO:15]

30

The hybridization reaction was carried out using  
equivalent amounts of each oligonucleotide (1 pm/μl final  
concentration for each oligo). The mixture of  
oligonucleotides was heated for 5 min at 100°C in ligation  
35 buffer (50 mM Tris-HCl, pH 7.5, 10 mM MgCl<sub>2</sub>, 10 mM DTT, 1mM

- 66 -

ATP, 25 µg/ml bovine serum albumin) and then cooled over at least 2 hours to 30°C.

The resulting hybridization product was ligated for 2 hours at room temperature or overnight at 16°C to the pJB02/Fc vector backbone which had been digested with *NheI* and *Eco47III*. The ligation products were used to transform competent XL-1 Blue cells (Stratagene). Recombinant plasmids were screened for the presence of peptide coding inserts by digesting clones with *NcoI* (encoding the Kozak sequence and first Met of the signal peptide) and sequenced. The resulting expression plasmid used for transfection assays was denoted pJB02-V8-GLP-1-Fc (Figure 5).

Example 1b Construction of DNA encoding Val<sup>8</sup>-GLP-1(7-37)-HSA: The plasmid HSA/pcDNA3.1GS was purchased from Invitrogen (Catalog # H-M12523M-pcDNA3.1/GS) and used as a template to isolate the cDNA encoding human serum albumin (HSA). The HSA cDNA was prepared using PCR wherein the DNA encoding the leader sequence as well as the six amino acid pro-peptide was removed from the 5' end. In addition, stop codons were added directly at the 3' end of the HSA coding sequence. Finally, restriction enzyme sites were engineered at the 5' and 3' end to facilitate cloning. The HSA DNA sequence present in the original vector purchased from Invitrogen contained a single base change in the 3' region of the gene (position 667) compared to the native human sequence. This change would result in a codon for Asn instead of Asp. Thus, using the strand overlapping PCR mutagenesis method discussed above, the codon was changed to code for Asp at this position. The resulting HSA encoding DNA was cloned into the *NheI* and *HindIII* sites of pJB02 to create pJB02-HSA (Figure 6).

The Igk leader sequence fused to the Val<sup>8</sup>-GLP-1(7-37) sequence was generated as discussed in Example 1a. This DNA

- 67 -

was ligated into the *NheI* and *FspI* sites of pJB02-HSA to create pJB02- Val<sup>8</sup>-GLP-1-HSA.

Example 1c Construction of DNA encoding Val<sup>8</sup>-GLP-1(7-37)-linker-HSA:

5 The vector pJB02-HSA was prepared as discussed in Example 1b. DNA encoding the linker sequence [GGGGG]<sub>3</sub> was ligated in frame to the 5' end of the HSA encoding DNA to create pJB02-linker-HSA (Figure 7). DNA encoding the Igk leader sequence and fused to the Val<sup>8</sup>-GLP-1(7-37) sequence 10 and the 5' part of the linker sequence was generated as discussed in Example 1a. This DNA was ligated into the *NheI* and *BspEI* sites of pJB02 to create pJB02- Val<sup>8</sup>-GLP-1-linker-HSA.

15

Example 1d Construction of DNA encoding Exendin-4-Fc:

The plasmid pJB02/Fc was prepared as described in Example 1a. DNA encoding the Igk signal sequence fused to Exending-4 was generated by *in vitro* hybridization of the 20 following overlapping and complementary oligonucleotides:

5' - CTAGCCACCATTGGAGACAGACACACTCCTGCTATGGGTACTGCTGCTCTG  
GGTTCAGGTTCCACCGGTCAC - 3' [SEQ ID NO:16]

25 5' - GGAGAGGGGAACCTTCACCGCGACCTGAGCAAGCAGATGGAGGAGGAGGCCGT  
GAGACTG - 3' [SEQ ID NO:17]

5' - TTCATCGAGTGGCTGAAGAACGGAGGACCAAGCAGCGGAGCCCCCTCCTCCT  
AGC - 3' [SEQ ID NO:18]

30

5' - GAACCTFGAACCAGAGCAGCAGTACCCATAGCAGGAGTGTGTCTGTCTCCA  
TGGTGG - 3' [SEQ ID NO:19]

35 5' - CTCCTCCTCCATCTGCTTGCTCAGGTCGCTGGTGAAGGTTCCCTCTCCGTGA  
CCGGTG - 3' [SEQ ID NO:20]



- 58 -

5' - GCTAGGAGGAGGGGCTCCGCTGCTTGGTCCTCCGTTCTTCAGCCACTCGAT  
GAACAGTCTTCACGSC - 3' [SEQ ID NO:21]

The hybridization reaction was carried out as described  
5 in Example 1a. The hybridized product was ligated to the  
pJB02 vector which had been digested with *NheI* and *Eco47III*  
as described in Example 1a to create pJB02-Exendin-4-Fc.

Example 1e Construction of DNA encoding Exendin-4-HSA:

10 The plasmid pJB02-HSA was prepared as described in  
Example 1b. DNA encoding the Igk signal sequence fused to  
Exending-4 was generated by *in vitro* hybridization of the  
same overlapping and complementary oligonucleotides  
described in Example 1d. Hybridization reactions were also  
15 carried out as described above. DNA was cloned into unique  
*NheI* and *FspI* sites in pJB02-HSA to create pJB02-Exendin-4-  
HSA.

20 Example 1f Construction of DNA encoding Exendin-4-linker-  
HSA:

The plasmid pJB02-linker-HSA was constructed as  
described in Example 1c. DNA encoding the Igk signal  
sequence fused to Exendin-4 and the 5' part of the linker  
sequence was generated as in Example 1d. This DNA was  
25 cloned into unique *NheI* and *BspEI* sites in pJB02-linker-HSA  
to create pJB02-Exendin-4-linker-HSA.

Example 1g Construction of DNA encoding Val<sup>8</sup>-GLP-1/C-Ex-Fc:

The plasmid pJB02-Exendin-4-Fc was prepared as  
30 described in Example 1d. The Exendin-4 encoding DNA was  
excised from the vector with *AgeI* and *Eco47III*. The Val<sup>8</sup>-  
GLP-1/C-Ex encoding DNA was generated by *in vitro*  
hybridization of the following overlapping and complementary  
oligonucleotides:

35

- 69 -

5' -CCGGTCACGTGGAGGGCACCTTCACCTCCGACGTGTCCTTCTATCTGGA  
GGGCCAGGCCGCCA - 3' [SEQ ID NO:22]

5' - AGGAATTCATCGCCTGGCTGGTGAAGGGCCGGGCAGCAGCGG  
5 AGCCCCCTCCTCCTAGC - 3' [SEQ ID NO:23]

5' - CTCCAGATAGGAGGACACGTCGGAGGTGAAGGTGCCCTCCAC  
GTGA - 3' [SEQ ID NO:24]

10 5' - GCTAGGAGGAGGGCTCCGCTGCTGCCCCGGCCCTTACCAGCCAGGCGA  
TGAATTCCTTGGCGGCTGGCC - 3' [SEQ ID NO:25]

The hybridization reaction was carried out as described  
in Example 1a. The hybridized product was ligated in place  
15 of Exendin-4 in the pJB02-Exendin-4-Fc expression vector to  
create pJB02-Val<sup>8</sup>-GLP-1/C-Ex-Fc.

Example 1h Construction of DNA encoding Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1-Fc:

The plasmid pJB02-Exendin-4-Fc was prepared as  
20 described in Example 1d. The Exendin-4 encoding DNA was  
excised from the vector with *AgeI* and *Eco47III*. The Val<sup>8</sup>-  
Glu<sup>22</sup>-GLP-1 encoding DNA was generated by *in vitro*  
hybridization of the following overlapping and complementary  
oligonucleotides:

25 5' -CCGGTCACGTGGAGGGCACCTTCACCTCCGACGTGTCCTTCTATCTCGA  
GGAGCAGGCCGCCA - 3' [SEQ ID NO:26]

5' - AGGAGTTCATCGCCTGGCTGGTGAAGGGCCGGGC - 3' [SEQ ID NO:27]

30 5' - GCCCCGGCCCTTACCAGCCAGGCGATGAACCTCTTGGCGGCC  
TGCTC - 3' [SEQ ID NO:28]

5' - CTCGAGATAGGAGGACACGTCGGAGGTGAAGGTGCCCT  
35 CCACGTGA - 3' [SEQ ID NO:29]

- 70 -

The hybridization reaction was carried out as described in Example 1a. The hybridized product was ligated in place of Exendin-4 in the pJB02-Exendin-4-Fc expression vector to create pJB02-Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1-Fc.

Example 1i Construction of DNA encoding Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1/C-Ex-Fc:

The plasmid pJB02-Exendin-4-Fc was prepared as described in Example 1d. The Exendin-4 encoding DNA was excised from the vector with *AgeI* and *Eco47III*. The Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1/C-Ex encoding DNA was generated by *in vitro* hybridization of the following overlapping and complementary oligonucleotides:

15 5' - CCGGTCACGTGGAGGGCACCTTCACCTCCGACGTGTCTCTCTATCTCGA  
GGAGCAGGCCGCCA - 3' [SEQ ID NO:30]

20 5' - AGGAATTCATCGCCTGGCTGGTGAAGGGCCGGGGCAGCAGCGGA  
GCCCCCTCTCTAGC - 3' [SEQ ID NO:31]

5' - CTCGAGATAGGAGGACACGTCGGAGGTGAAGGTGCC  
TCCACGTGA - 3' [SEQ ID NO:32]

25 5' - GCTAGGAGGAGGGGCTCCGCTGCTGCCCGGCCCTTCACCAGCCAGGCCA  
TGAATTCCTTGGCGGCTGCTC - 3' [SEQ ID NO:33]

The hybridization reaction was carried out as described in Example 1a. The hybridized product was ligated in place of Exendin-4 in the pJB02-Exendin-4-Fc expression vector to create pJB02-Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1/C-Ex-Fc.

Example 1j Construction of DNA encoding Gly<sup>8</sup>-GLP-1-Fc:

The plasmid pJB02-Exendin-4-Fc was prepared as described in Example 1d. The Exendin-4 encoding DNA was

- 71 -

excised from the vector with AgeI and Eco47III. The Gly<sup>8</sup>-GLP-1 encoding DNA was generated by *in vitro* hybridization of the following overlapping and complementary oligonucleotides:

5

5' - CCGGTCACGGCGAGGGCACCTTCAGTACGTGTCTCTCTATCTGGA  
GGGCCAGGCCGCCA - 3' [SEQ ID NO:34]

10

5' - AGGAGTTCATCGCCTGGCTGGTGAAGGGCCGGGGC - 3' [SEQ ID NO:35]

5' - CTCACAGATAGGAGGACACGTCAGTGAAGGTGCCCTC  
GCCGTGA - 3' [SEQ ID NO:36]

15

5' - GCCCCGGCCCTTCACCAGCCAGGCGATGAACCTCTGGCGGC  
CTGGCC - 3' [SEQ ID NO:37]

The hybridization reaction was carried out as described in Example 1a. The hybridized product was ligated in place of Exendin-4 in the pJB02-Exendin-4-Fc expression vector to  
20 create pJB02-Gly<sup>8</sup>-GLP-1-Fc.

#### Example 2: Expression of heterologous fusion proteins

Expression of the fusion proteins encoded by the DNA constructs of Example 1 was carried out by transiently  
25 transfecting HEK 293BNA cells (both adherent and suspension). Cells were counted and seeded 24 hours prior to transfection. The transfection cocktail was prepared by mixing FuGene<sup>TM</sup>6 transfection reagent (Roche Molecular Biochemicals, catalog # 1814443) with OptiMEM (Gibco/BRL)  
30 and incubating at room temperature for 5 min at which point DNA was added and the cocktail was incubated for an additional 15 min. Immediately before transfection, fresh growth media was added to the plate. Tables 1 and 2 provide further transfection details.

35

- 72 -

Table 1: Reagents used in transient transfections of 293EBNA cells.

Tissue culture vessel	Number of cells seeded	DNA ( $\mu$ g)	FuGene ( $\mu$ l)	OptiMEM media (ml)	Vol. of growth medium (ml)
35 mm	$5 \times 10^5$	1.5	9	0.102	2
100 mm	$2 \times 10^6$	12	73	0.820	10
700 cm <sup>2</sup> (RB)	$2 \times 10^7$	65	400	4.0	100

5 Table 2: Media composition

Growth and transfection medium	Harvesting medium
DMEM F12 3:1	Hybritech base
5 % FBS	1 mM Ca <sup>2+</sup>
20 mM HEPES	20 mM HEPES
2 mM L-glutamine	1 $\mu$ g/ml Nuselin (human insulin)
50 $\mu$ g/ml geneticin (G418 NEO)	1 $\mu$ g/ml human transferrin
50 $\mu$ g/ml tobromycin	50 $\mu$ g/ml tobromycin

For small-scale transfections (35mm - 10mm vessels), cells were rinsed with PBS and switched to harvesting media 24 hours post- transfection and media was collected and replaced every 24 hours for several days. In the case of large-scale transfections (700 cm<sup>2</sup> roller bottles), the roller bottles were rinsed with PBS 48 hours post-transfection and changed to harvesting media. Media was collected and changed every 24 hours for at least 10 consecutive days. Routinely, only 10 harvests were used for subsequent protein purification.

- 73 -

Example 3: Purification of heterologous fusion proteinsExample 3a purification of Val<sup>8</sup>-GLP-1-Fc

Approximately 4.5 liters of conditioned medium (fusion protein expression level approximately 20 µg/ml) from large-scale transfections was filtered using a CUNO filter system and concentrated to 250 ml using a ProFlux tangential flow filtration system with a 10 K filter membrane. Val<sup>8</sup>-GLP-1-Fc was captured with a 5 ml HiTrap protein A column in 1x PBS, pH 7.4 at a flow rate of 2 ml/min and eluted with 50 mM citric acid pH 3.3. Fractions (1 ml) were collected in tubes containing 4 ml 1x PBS and 100µl 1 M Tris pH 8.

Fractions containing the fusion protein, as determined by SDS-PAGE and reverse phase-HPLC on Zorbax C8, were pooled and applied to a Superdex 75 60/60 column in 1x PBS pH 7.4 at a flow rate of 10 ml/min. Positive fractions (20 mls/tube) were collected and pooled. Pooled fractions were then subjected to C4 Reverse Phase Chromatography in 0.1 %TFA water at a flow rate of 3 ml/min. Val<sup>8</sup>-GLP-1-Fc was eluted using a gradient from 5% B (0.1% TFA in acetonitrile) to 100% B in 70 min. Eluant fractions (3 mls/tube) were collected. Acetonitrile was removed by vacuum drying and 1 ml of H<sub>2</sub>O was added. The purified sample (approximately 32 mls) was dialyzed twice against 4 liters of 1x PBS pH7.4.

The dialyzed sample was then filtered using a MILLEX-GV 0.22 µm Filter Unit and concentration was determined using absorption at 280 nm.

Example 3b purification of Val<sup>8</sup>-GLP-1-HSA or Val<sup>8</sup>-GLP-1-Linker-HSA

Approximately 5.5 liters of conditioned medium (fusion protein expression level approximately 10µg/ml) was filtered using a CUNO filter system and concentrated to 380 mls using a ProFlux tangential flow filtration system with a 10 K filter membrane.

- 74 -

The fusion protein was captured using a 50 ml Fast Flow Q column (Pharmacia) in 20 mM Tris pH 7.4 at a flow rate of 5ml/min. Protein was eluted using a gradient: from 0% to 50% 20mM Tris pH 7.4, 1M NaCl in 10 CV, then to 100%B in 2 CV.

5 Fractions containing the fusion protein were pooled and subjected to C4 Reverse Phase Chromatography in 0.1% TFA water at a flow rate of 5 ml/min. The fusion protein was eluted using a gradient from 20% B (0.1% TFA in acetonitrile) to 90% B in 120min. Fractions (3.5 ml/tube)  
10 were collected. Acetonitrile was removed by vacuum drying.

Approximately 9 mls of pooled sample was diluted with 1x PBS pH 7.4 to 40ml and dialyzed against 4 liters of 1x PBS pH 7.4 overnight. The sample was filtered and concentration was determined by absorption at 280nm.

15

Example 3c purification of Exendin-4-Fc:

Approximately 4 liters of conditioned medium (fusion protein expression level approximately 8 µg/ml) was filtered using a CUNO filter system and concentrated to 250 mls using  
20 a ProFlux tangential flow filtration system with a 30K filter membrane.

Exendin-4-Fc was captured with a 5 ml HiTrap protein A column in 1x PBS, pH 7.4 at a flow rate of 2 ml/min and eluted with 50 mM citric acid pH 3.3. Fractions containing  
25 the fusion protein were pooled, filtered, and dialyzed against 4 liters of 1 x PBS over night. The dialyzed sample was then applied to a Superdex 75 60/60 column in 1x PBS pH7.4, 0.5M NaCl at a flow rate of 10 ml/min. Fractions (20 ml/tube) containing the fusion protein were collected,  
30 pooled, and concentrated to about 1 mg/ml. Concentrated samples were then filtered using a MILLEX-GV 0.22 µm Filter Unit.

- 75 -

Example 3d purification of Exendin-4-HSA and Exendin-4-linker-HSA:

Approximately 1.1 liters of conditioned medium (fusion protein expression level approximately 6 $\mu$ g/ml) was filtered using a CUNO filter system and concentrated to 175 mls using a ProFlux tangential flow filtration system with a 30K filter membrane.

The fusion protein was captured using a 5 ml HiTrap Q-sepharose column (Pharmacia) in 20 mM Tris pH 7.4 at a flow rate of 2 ml/min. Protein was eluted using a gradient from 0% to 50% 20mM Tris pH 7.4, 1M NaCl in 12 CV and then to 100%B in 4 CV.

Fractions containing the fusion protein were pooled and subjected to C4 Reverse Phase Chromatography in 0.1% TFA water at a flow rate of 5 ml/min. The fusion protein was eluted using a gradient from 10% B (0.1% TFA in acetonitrile) to 100% B in 70 min. Fractions (10 ml/tube) containing the fusion protein were collected. Acetonitrile was removed using a vacuum dryer.

Approximately 8 mls of pooled sample was dialyzed against 4 liters of 1x PBS pH 7.4 overnight. The sample was filtered and concentration was determined by absorption at 280nm. The dialyzed sample was then applied to a Superdex 200 26/60 column in 1x PBS pH 7.4, 0.5 M NaCl at a flow rate of 2 ml/min. Fractions (3 ml/tube) containing the fusion protein were collected, pooled, concentrated, and filtered.

Example 4: Characterization of fusion proteins by SDS PAGE:

SDS-PAGE followed by immunoblotting was used to analyze both purified fusion protein as well as conditioned medium from cells transfected with various fusion protein expression vectors. SDS-PAGE was performed on a Novex Powerease 500 system using Novex 16% Tris-Glycine Precast gels (EC6498), running buffer (10x, LC2675) and sample



- 76 -

buffer (L2676). Samples were reduced with 50 mM DTT and heated 3-5 min at 95°C prior to loading.

After running the SDS-PAGE gel, water and transfer buffer (1X Tris-Glycine Seprabuff (Owl Scientific Cat. No. ER26-S) with 20% methanol) were used to rinse SDS from the gels. A Novex transfer apparatus was used with PVDF (BioRad, Cat. No. 162-0174) and nitrocellulose membranes (BioRad, Cat. No. 1703965 or 1703932). Transfer was carried out at room temperature for 90 min at 30-35 V. Membranes were  
10 blocked in 1X PBS with 0.1% Tween-20 (Sigma, Cat. No. P-7949) and 5% Milk (BioRad, Cat. No. 170-6404) for 1-12 hours at 4°C. Antibodies are diluted into 1X PBS +5% Milk and the blots are incubated in these solutions for 1-2 h at 4°C. Between incubations, the blots are washed four times for 5  
15 min each with 1X PBS and 0.2% Tween-20 at room temperature. PBS was made from either GIBCO 10X PBS (Cat No. 70011), to give a final composition of 1 mM monobasic potassium phosphate, 3 mM dibasic sodium phosphate, 153 mM sodium chloride, pH 7.4, or PBS pouches from Sigma (Cat. No. 1000-  
20 3), to give 120 mM NaCl, 2.7 mM KCl and 10 mM phosphate, pH 7.4 at 25°C.

The primary antibodies were either a polyclonal goat anti-IgG1 or rabbit anti-HSA. The secondary antibody was either an anti-goat IgG HRP or an anti-rabbit IgG HRP. The  
25 secondary antibody was diluted 1:5000. An ECL system (Amersham Pharmacia Biotech, Cat. No. RN2108 and Cat. No. RPN1674H) was used for developing blots.

Figure 3A compares purified Fc protein to conditioned media from pJB02-Val<sup>8</sup>-GLP-1-Fc and pJB02-Exendin-4-Fc  
30 transfected cells. The decrease in mobility is consistent with the increased size due to the GLP-1 portion of the fusion protein. Figure 3B similarly compares purified HSA with conditioned media from cells transfected with pJB02-Val<sup>8</sup>-GLP-2-HSA, pJB02-Val<sup>8</sup>-GLP-1-Linker-HSA, pJB02-Exendin-

- 77 -

4-HSA, or pJB02-Exendin-4-Linker-HSA. Figure 4 identifies purified fusion protein preparations.

Example 5: Characterization of fusion  
proteins using mass spectrometry:

5 All experiments were performed on a Micromass ToFSpec-2E mass spectrometer equipped with Time Lag Focusing electronics, a Reflectron (used in analysis of the 0-8000 Da peptide range), a Linear detector (used during high mass / good signal analysis), and Post Acceleration Detector (or P.A.D., used for high mass / extremely low signal analysis). The effective path length of the instrument in Linear mode is 1.2 meters, in Reflectron mode it is 2.3 meters. Two dual micro-channel plate detectors are fitted for linear and reflectron mode detection. The laser used is a Laser Science Inc. VSL-337i nitrogen laser operating at 337 nm at 15 laser shots per second. All data were acquired using a 2 GHz, 8 bit internal digitizer and up to 50 laser shots were averaged per spectrum.

20 The instrument was operated in linear mode for the analysis of the GLP-1 fusion proteins in question. The linear detector is a device that detects ions that travel down the flight tube of the MALDI-ToF-MS instrument. It measures the ion abundance over time and sends a signal to the digitizer for conversion. The digitizer is an analog-to-digital converter that allows the signal from the mass spectrometer to be transferred to the computer, where it is reconstructed into a usable m/z spectrum.

25 A recrystallized saturated sinapinic acid solution (diluted in 50/50 Acn / H<sub>2</sub>O and 0.1% TFA) was utilized as the ionization matrix. Sinapinic acid is a proper matrix for proteins above 10 kDa. Mass appropriate reference proteins were used for internal and external calibration files in order to obtain accurate mass determinations for the samples 30 analyzed. Samples were all analyzed using a 1:2 sample to

- 78 -

matrix dilution. The instrument was initially set up under the following linear detector conditions:

Source Voltage: 20.0 keV      Pulse Voltage: 3.0 keV  
 5 Extraction Voltage: 20.0 keV      Laser Coarse: 50  
 Focus Voltage: 16.0 keV      Laser Fine: 50  
 Linear detector: 3.7 keV  
 P.A.D.: (off line)

10 These settings were modified (as needed) to give the best signal/noise ratio and highest resolution. Table 3 provides a characterization of different GLP-1 fusion proteins.

15 **Table 3**

Fusion Protein	Expected Mass (KDa)	Mass determined by Mass Spec (kDa)
Val <sup>8</sup> -GLP-1-IgG1	59.08	61.94
Val <sup>8</sup> -Glu <sup>22</sup> -GLP-1-IgG1	59.23	63.61
Gly <sup>8</sup> -GLP-1-IgG1	59.00	62.93
Val <sup>8</sup> -GLP-1-CEX-IgG1	60.45	65.1-65.6
Val <sup>8</sup> -Glu <sup>22</sup> -GLP-1-CEX-IgG1	60.69	65.86
Exendin-4-IgG1	60.69	65.86
Val <sup>8</sup> -GLP-1-Linker-HSA	70.70	69.89, 70.74
Exendin-4-HSA	70.56	70.62
Exendin-4-Linker-HSA	71.56	71.62

CEX refers to a C-terminal extension and comprises the sequence of Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser.

Linker is Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly- Gly-Gly-Gly-Ser.

20

- 79 -

Example 6: Activity of heterologous fusion proteins:

The ability of the fusion proteins of the present invention to activate the GLP-1 receptor was assessed using in vitro assays such as those described in EP 619,322 to Gelfand, et al., and U.S. Patent No. 5,120,712, respectively. The activity of these compounds relative to the activity of Val<sup>8</sup>-GLP-1(7-37)OH is reported in Table 4. Figure 8 represents in vitro dose response curves for Val<sup>8</sup>-GLP-1 and Exendin-4 fusion proteins. In addition, Table 5a and 5b provide the in vitro activity of a large group of GLP-1 analogs that can be fused to an Fc or an albumin protein to make biologically active fusion proteins. These activities are compared to GLP-1(7-37)OH.

15 Table 4: In vitro activity of GLP-1 fusion proteins

Fusion Protein	In Vitro Activity (% of Val <sup>8</sup> -GLP-1)
Val <sup>8</sup> -GLP-1-IgG1	1
Exendin-4-IgG1	240
Val <sup>8</sup> -GLP-1-Linker-HSA	0.2
Exendin-4-HSA	20
Exendin-4-Linker-HSA	90
Exendin-4	500
Val <sup>8</sup> -Glu <sup>22</sup> -GLP-1-IgG1	3.7
Gly <sup>8</sup> -GLP-1-IgG1	3.3
Val <sup>8</sup> -GLP-1-CEX-IgG1	3.3
Val <sup>8</sup> -Glu <sup>22</sup> -GLP-1-CEX-IgG1	29
Gly <sup>8</sup> -Glu <sup>22</sup> -GLP-1-C2-IgG1	75
Gly <sup>8</sup> -Glu <sup>22</sup> -GLP-1-CEX-Linker-IgG1	150
Exendin-4-C2-IgG1	250
Exendin-4-Linker-IgG1	330
Gly <sup>8</sup> -Glu <sup>22</sup> -GLP-1-CEX-Linker-HSA	4
Gly <sup>8</sup> -Glu <sup>22</sup> -GLP-1-CEX-Linker-IgG4	80

- 80 -

CEx refers to a C-terminal extension and comprises the sequence of Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser. Linker is Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Ser

- 5 C2 is Ser-Ser-Gly-Ala-Ser-Ser-Gly-Ala.

The amino acid sequences of the fusion proteins described in Tables 3 and 4 are represented in SEQ ID NO: 13 to SEQ ID NO: 31.

Val<sup>8</sup>-GLP-1-Human serum albumin amino acid sequence is

- 10 represented by SEQ ID NO: 13.

1 HVEGTFTSDV SSYLEGQAAK EPIAWLVKGR GDAHKSEVAH RFDLGEZNF KALVLIAFAQ  
 61 YLQQCFPEDH VKLVNEVTEF ARTCVADESA ENCDKSLHTL PGDKLCTVAT LRETYGEMAD  
 121 CCAKQEPERN ECPQLHKDDN PNLKPLVRPE VDVKTAFPHD NBETFLKKYL YEIAKRHPYF  
 181 YAPELLFFAK RYKAAFTGCC QAADKAACIL PKLDELRLDEG KASSAKRLK CASLQKFGER  
 15 241 AFKANAVARL SQRFPAEFA EVSKLVTDLT KVHTECCRGD LLECAADDRAD LAKYLCEQND  
 301 SISSEKKECC EKPLLEKSHC IAEVENDEMP ADLPSLAADF VESKDVCKNY AEAKDVFLGM  
 361 FLYEYARRHP DYSVVLLRL ARTYETTFLEK CCAAADPHEC YAKVFDEFKP LVEEPQNLK  
 421 QNCELFEQLG EYKFNALLV RYTKKVPQVS TPTLVEVSRN LGKVGSKCK HPAKRMECA  
 481 EDYLSVVLNQ LCVLHEKTPV SDRVTKCCTE SLVNRRCPCS ALEVDETYVE KFNATETTF  
 20 541 HADICTLSEK ERQIKKQAL VELVKKHKA TRQLKAVMD DFAAPVEKCC KADDKETCPA  
 601 BEGKKLVAAS QAALGL [SEQ ID NO: 13]

Val<sup>8</sup>-GLP-1-Linker-Human serum albumin amino acid sequence is represented by SEQ ID NO: 14.

25 1 HVEGTFTSDV SSYLEGQAAK EPIAWLVKGR GGGGGSGGGG SGGGGSDAHK SEVANHFKDL  
 61 GSENFKALVL IAFAYLQCC PFEDHVKLVN EVTEPAKTCV ADESAENCDK SLHTLPGDKL  
121 CTVTATLRETY GEMADCCAKO EPERNECFLO HKDDNFNLPR LVRPEVDVMC TAFHNBETTF  
181 LKKYLYELAR RHPYFYAPEL LFFAKRYKAA FTECCQAADK AACLLPKLDE LRDEGKASSA  
 241 KQSLKCAQLG KFGERAFKAN AVARLSQRF EKAFAEVS KLVTOLTKVHTE CCGDLLECA  
 30 301 DDRADLAKYI CENQDSISS LKECCCKPLL EKSHCIAEVE NDEMPADLPS LAADPVESKD  
 361 VCKNYAEAK VFLGMPLEYE ARRHFDYSVV LLLRLAKTYE TILSKCCAAA DFHECYAKVF  
 421 DEFKPLVEEP QNLKQNCLE FBQLGEYKFO NALLVRYTKK VFQVSTPTLV EVSRNLKRGV  
 481 SKCCKHPEAK KMPCAEDYLS VVLNQLCVLH EKTVPVSDRV KCTPESLVNR PCFSALEVD  
 541 EYTVKPEFNA ETTFPHADIC TLSEKERQIK KQTLVVELVK HKPKATKEQL KAVMDPFAAF  
 35 601 VRKCKADK ETCFAEGKK LVAASQAALG L [SEQ ID NO: 14]

- 81 -

Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-Human serum albumin amino acid

sequence is represented by SEQ ID NO: 15.

```

1 HGEGTPTSDV SSYLEBQAAK EFLAWLVKGR GSSGAPPPSG GGGGSGGGGS GGGGSDAHKS
61 EVAHRPKDLG EENFKALVLI APAQYLQQCP FEDHVKLVNE VTEFAKTCVA DESAENCCKS
5 121 LHTLFGDKLC TVATLRBTYG EMADCCAKQE PERNECFLOH KDDNPNLPRL VREFVDVMCT
181 AFPHNRETFI KKYLYEIAIR HPYFYAPELL FFAKRYKAAF TECCQAADKA ACLLPKLEDEL
241 RDEGKASSAK QRLKASLQK FGERAFKAWA VARLSQRFPK AEFAEVSKLV TDLTVHTEC
301 CHGDLEECAD DRADIAKYIC ENQDSISSKL KECCKELPKE KSHCIAEVEN DEMPADLPESL
361 AADPVESKDV CKNYAEAKDV FLGMFLYHYA RRPDPYSVVL LLRLAKTYET TLEKCCAAAD
10 421 PHECYAKVPD EPKPLVREFQ NLIKQNCLEF EQLGEYKQFN ALLVRYTKKV PQVSTPTLVE
481 VSRNLGKVGK KCKKHPEAKR MPCAEDYLSV VLNQLCVLHE KTFVSDRVTK CCTESLVNRR
541 PCFSALEVDE TYVPKEFMAS TTFPHADICT LSEKERQIKK QTALVELVKH KPAKTEQILK
601 AVMDFAAFV EKCKKADDEK TCPABEGKKL VAASQALGL [SEQ ID NO: 15]

```

15 Exendin-4-Human serum albumin amino acid sequence is represented by SEQ ID NO: 16.

```

1 HGEGTPTSDL SKMEEBAVR LFIEWLKNGG PSSGAPPPSD AHKSEVAHRF KDLGEENPKA
61 LVLIAPAQHL QQCPFEDHVK LVNEVTEFAK TCVADESAEN COKSLHTLFG DKLCIVATLR
121 ETYGEMADUC AKQEPERNEC FLQHKDDNPN LPRLVRPEVD VMCTAFHDNE ETFLKKLYE
20 181 IARRHPYFYA PELLFFAKRY KAAFTCCQA ADKAACLLEK LDELRLDEGKA SSARKRLKCA
241 SLQKFGGERAF KAWAVARLSQ RFPKAEFAEV SKLVTDLTKV HTECCGDEL ECADDRADYA
301 KYICENQDSI SSKLECCCEK PILEKSHCIA EVENDENPAD LPSLAADFVE SKDVCKNYAE
361 AKDVFLGMFL YEYARRHPDY SVVILLRLAK TYETTLKCKC AAADPHECTA KVPDEFKPLV
421 KEPQNLIKQN CELFBQLGEY KQNALLVRY TKKVPQVSTP TLVEVSRNLG KVGSKCKKHP
25 481 BAKRMPCAED YLSVVLNQLC VLHKEFVSD RVTECCTESL VNRNPFCSAL EVDETYVPKE
541 FNAETFTFFH DICTLSRER QIRKQTALVE LVKHKPKATK EQLKAVMDFF AAFVEKCKCA
601 DDKETCPASE GKILVAASQA ALGL [SEQ ID NO: 16]

```

30 Exendin-4-Linker-Human serum albumin amino acid sequence is represented by SEQ ID NO: 17.

```

1 HGEGTPTSDL SKMEEBAVR LFIEWLKNGG PSSGAPPPSG GGGGSGGGGS GGGGSDAHKS
61 EVAHRPKDLG EENFKALVLI APAQYLQQCP FEDHVKLVNE VTEFAKTCVA DESAENCCKS
121 LHTLFGDKLC TVATLRBTYG EMADCCAKQE PERNECFLOH KDDNPNLPRL VREFVDVMCT
181 AFPHNRETFI KKYLYEIAIR HPYFYAPELL FFAKRYKAAF TECCQAADKA ACLLPKLEDEL
35 241 RDEGKASSAK QRLKASLQK FGERAFKAWA VARLSQRFPK AEFAEVSKLV TDLTVHTEC
301 CHGDLEECAD DRADIAKYIC ENQDSISSKL KECCKELPKE KSHCIAEVEN DEMPADLPESL
361 AADPVESKDV CKNYAEAKDV FLGMFLYHYA RRPDPYSVVL LLRLAKTYET TLEKCCAAAD
421 PHECYAKVPD EPKPLVREFQ NLIKQNCLEF EQLGEYKQFN ALLVRYTKKV PQVSTPTLVE

```

- 82 -

481 VSRNLGKVGSG KCKKHPEAKR MPCAEDYLSV VLNQLCVLHSE KTPVSDRVTK COTESLVNRR  
 541 PCFSALEVDSE TVVPKEFNAB TPTPHADICT LSEKERQIKK QTALVELVKH KPKATKESQLK  
 601 AVMDDPFAAFV ERCKKADDKE TCFARHGKSL VAASQAALGL [SEQ ID NO: 17]

5 Val<sup>8</sup> --GLP-1-IgG1 amino acid sequences represented by SEQ ID NO: 18.

1 HVEGTFTSDV SSYLEGQAAK EPIANLVKGR GAEPKSCDKT HTCPPCPAPE ILGGPSVFLF  
 61 PPKPKDTLMI SRTPVETCVV VDVSHEDPEV KFNWYVDGVE VHNAKTKPRE EQYNSTYRVV  
 121 SVLTVLHQDN LMGKEYKCKV SNKALPAPIE KTISKAKGQP REPQVYTLPP SREMTKNQV  
 10 181 SLTCLVKGFY PSDIAVWES NGQPENNYKT TFPVLDSGGS FFLYSLKLTVD KSRWQQGNVF  
 241 SCSVMHEALH NHYTQKSLSL SPGK [SEQ ID NO: 18]

Val<sup>8</sup> --GLP-1-Cex-IgG1 amino acid sequence is represented by SEQ ID NO: 19.

15 1 HVEGTFTSDV SSYLEGQAAK EPIANLVKGR GSSGAPPPSA EPKSCDKTHT CPFCPAPELL  
 61 GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NMYVDGVEVH NAKTKPREEQ  
 121 YNSTYRVVSV LTVLHQDNLN GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR  
 181 REMTKNQVSL TCLVKGFPYS DIAVWESNG QPENNYKTFP VLDSGGSFF LYSKLTVDKS  
 241 RWQQGNVFSC SVMHEALHNN YTQKSLSLSP GK [SEQ ID NO: 19]

20

Val<sup>8</sup> --Glu<sup>22</sup> GLP-1-IgG1 amino acid sequence is represented by SEQ ID NO: 20.

1 HVEGTFTSDV SSYLEGQAAK EPIANLVKGR GAEPKSCDKT HTCPPCPAPE LLOGPSVFLF  
 61 PPKPKDTLMI SRTPVETCVV VDVSHEDPEV KFNWYVDGVE VHNAKTKPRE EQYNSTYRVV  
 25 121 SVLTVLHQDN LMGKEYKCKV SNKALPAPIE KTISKAKGQP REPQVYTLPP SREMTKNQV  
 181 SLTCLVKGFY PSDIAVWES NGQPENNYKT TFPVLDSGGS FFLYSLKLTVD KSRWQQGNVF  
 241 SCSVMHEALH NHYTQKSLSL SPGK [SEQ ID NO: 20]

Val<sup>8</sup> --Glu<sup>22</sup> GLP-1-CEx-IgG1 amino acid sequence is represented by SEQ ID NO: 21.

30 1 HVEGTFTSDV SSYLEGQAAK EPIANLVKGR GSSGAPPPSA EPKSCDKTHT CPFCPAPELL  
 61 GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NMYVDGVEVH NAKTKPREEQ  
 121 YNSTYRVVSV LTVLHQDNLN GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR  
 181 EEMTKNQVSL TCLVKGFPYS DIAVWESNG QPENNYKTFP VLDSGGSFF LYSKLTVDKS  
 35 241 RWQQGNVFSC SVMHEALHNN YTQKSLSLSP GK [SEQ ID NO: 21]

- 83 -

Gly<sup>8</sup>-Glu<sup>22</sup> GLP-1-C2-IgG1 amino acid sequence is represented by SEQ ID NO: 22.

1 HGEGTFTSDV SSYLEEQAAR EPIANLVKGR GSSGASSGAA EPKSCDKTHT CPFCFAPELL  
61 GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NNYVDGVEVH NAKTKPREEQ  
5 121 YNSTYRVVSV LTVLHQDWLN GKEYKCKVSN KALPAPIEKT ISKAKQPRE PQVYTLPPSR  
181 EEMTKNQVSL TCVLKGFYPS DIAVEWESNG QPENNYKTFP PVLDSGGSFF LYSKLTVDKS  
241 RWQQGNVFSC SVMHEALHNN YTQKSLSLSP GK [SEQ ID NO: 22]

Gly<sup>8</sup>-Glu<sup>22</sup> GLP-1-CEx-Linker-IgG1 amino acid sequence is represented by SEQ ID NO: 23.

1 HGEGTFTSDV SSYLEEQAAR EPIANLVKGR GSSGAPPPSG GGSGGGGSG GGSAAEPKSC  
61 DKHTCTPFCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNMYVD  
121 GVEVHNAKTK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK  
181 QQPREPQVYT LPSPREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQFENN YKTFPPVLDG  
15 241 DGSFFLYSKL TVDKSRWQQG NVFSCSVMEH ALNHYHTQKS LSLSPGK [SEQ ID NO: 23]

Gly<sup>8</sup>-Glu<sup>22</sup> GLP-1-CEx-Linker-IgG4 amino acid sequence is represented by SEQ ID NO: 24.

1 HGEGTFTSDV SSYLEEQAAR EPIANLVKGR GSSGAPPPSG GGSGGGGSG GGSAAESKYG  
20 61 PFCSPSCPAFE FLGGPSVFLF FPKPKDTLMI SRTPEVTCVV VDVSGEDPEV QFMWYVDGVE  
121 VHNAKTKPRE EQFNSTYRVV SVLTVLHQDW LNKKEYKCKV SNKGLPSSIE KTKISKAKGP  
181 RFEPQVYTLPP SQEEMTKNQV SLTCLVKGFY PSDIAVEWES NGQPENNYKT TPVLDSDGGS  
241 FFLYSRLTVD KSRMQEGNVF SCSVMHEALH NHYTKLSLSL SLGK [SEQ ID NO: 24]

Gly<sup>8</sup>-Glu<sup>22</sup> GLP-1-CEx-2Linker-IgG1 amino acid sequence is represented by SEQ ID NO: 25.

1 HGEGTFTSDV SSYLEEQAAR EPIANLVKGR GSSGAPPPSG GGSGGGGSG GGSGGGGSG  
61 GGSGGGGGSA EPKSCDKTHT CPFCFAPELL GGPSVFLFPP KPKDTLMISR TPEVTCVVVD  
121 VSHEDPEVKF NNYVDGVEVH NAKTKPREEQ YNSTYRVVSV LTVLHQDWLN GKEYKCKVSN  
30 181 KALPAPIEKT ISKAKQPRE PQVYTLPPSR EEMTKNQVSL TCVLKGFYPS DIAVEWESNG  
241 QPENNYKTFP PVLDSGGSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNN YTQKSLSLSP  
301 GK [SEQ ID NO: 25]

Gly<sup>8</sup>-Glu<sup>22</sup> GLP-1-2Linker-IgG1 amino acid sequence is represented by SEQ ID NO: 26.



- 84 -

1 HGEGETFTSDV SSYLEEQAAK EPIAWLVKGR GGGGSGGGG SGGGSGGGG SGGGSGGGG  
 61 SAEPKSCDKT HTCPPCPAP E LGGPSVFLF PKPKDTLMI SRTPEVTCVV VVVSHPEDPEV  
 121 KFNWYVDGVE VHNKTKPRE EQYNSTYRVV SVLTVLHQDW LNKKEYKCKV SNKALPAPIE  
 181 KTISKAKGQP REFQVYTLPP SREEMTKNQV SLTCLVKGfy PSDIAVENES NQQPENNYKT  
 241 TFPVLDSGGS FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL SPGK  
 5 [SEQ ID NO: 26]

Gly<sup>8</sup>-Glu<sup>22</sup> GLP-1-2CEX-IgG1 amino acid sequence is represented  
 by SEQ ID NO: 27.

10 1 HGEGETFTSDV SSYLEEQAAK EPIAWLVKGR GSSGAPPPSS SGAPPPSAEP KSCDKTHTCP  
 61 PCPAPELLGG PSVFLPPPKP KDTLMISRTP EVTCVVVDVS HEDFPEVKFNW YVDGVEVHNA  
 121 KTKPREEQYN STYRVVSVLT VLHQDWLNKG EYCKCKVSNKA LPAPIEKTIS KAKGQPREPQ  
 181 VYTLPPSREE MTKNQVSLTC LVKGFYPSDI AVENESNGQP ENNYKTPPV LDSGGSFFLY  
 241 SKLTVDKSRW QQGNVFSVCSV MHEALHNHYT QKSLSLSPGK [SEQ ID NO: 27]

15

Gly<sup>8</sup>-Glu<sup>22</sup>-Val<sup>25</sup>-Ile<sup>33</sup> GLP-1-CEX-Linker-IgG1 amino acid  
 sequence is represented by SEQ ID NO: 28.

1 HGEGETFTSDV SSYLEEQAAK EPIAWLVKGR GSSGAPPPSS GGGSGGGGSG GGSAAEPKSC  
 61 DKTHTCPPCP APPELLGGPSV FLFPPPKPDT LMISRTPEVT CUVVDVSHED PEVKFNWYVD  
 20 121 GVEVHNKATK PREEQYNSTY RVVSVLTVLH QDWLNKEYK CKVSNKALPA PIEKTISKAR  
 181 GQPREPQVYT LPFSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPEVLDS  
 241 DGSFFLYSKL TVDKSRWQQG NVFSCSVME ALHNHYTQKS LSLSPGK [SEQ ID NO: 28]

Exendin-4-IgG1 amino acid sequence is represented by SEQ ID  
 25 NO: 29.

1 HGEGETFTSDL SKQMEEEAVR LFIEMLNKG PSSGAPPPSA EPKSCDKTHT CPPCPAPELL  
 61 GGPVSFLPPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NQWYVDGVEVH NAKTKPREQ  
 121 YNSTYRVVSV LTVLHQDWLN GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR  
 181 EEMTKNQVSL TCLVKGfyps DIAVENESNG QPENNYKITE PVLDSGGSFF LYSKLTVDKRS  
 30 241 NQQQGNVFS CVMHEALHNH YTKRSLSLSP GK [SEQ ID NO: 29]

- 85 -

Exendin-4-C2-IgG1 amino acid sequence is represented by SEQ ID NO: 30.

1 HGEETPTSDL SKQMEERAVR LFIEWLKNKG PSSGASSGAA EPKSCDKTHT CPFCPAPELL  
 5 61 GGPSVFLPPP KPKDILMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVR NAKTKPREEQ  
 121 YNSTYRVVSV LTVLHQDWLN GKEYKCKVSN KALPAPIEKT ISKAKQPRE PQVYTLPPSR  
 181 EEMTKNQVSL TCLVKGFPYS DIAVEWESNG QPENNYKTF PVLDSGGSFF LYSKLTVDKS  
 241 RWQQGNVFSC SVMHEALHNN YTQKSLSLSP GK [SEQ ID NO: 30]

10 Exendin-4-Linker-IgG1 amino acid sequence is represent by SEQ ID NO: 31.

1 HGEETPTSDL SKQMEERAVR LFIEWLKNKG PSSGAPPSG GGGSGGGSG GGGSAEPKSC  
 61 DKIRTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD  
 121 GVEVHNATK PREEQYNSTY RVVSVLTVLN QDWLNGKEYK CKVSNKALPA PIEKTISKAK  
 15 181 QQPREPQVYT LPFSRREMTK NQVSLTCLVK GPYPSDIAVE WESRGQPENN YKITPPVLD  
 241 DGSFFLYSKL TVDKSRWQQG NVFSCSVME ALHNHYTQKS LSLSPGK [SEQ ID NO: 31]

Table 5a: *In vitro* GLP-1 analog activity

GLP-1 Receptor	
<u>GLP-1 Compound</u>	<u>Activation</u>
20 GLP-1(7-37)OH	1.0
Val <sup>8</sup> -GLP-1(7-37)OH	0.47
	(n = 6)
25 Gly <sup>8</sup> -His <sup>11</sup> -GLP-1(7-37)OH	0.282
Val <sup>8</sup> -Ala <sup>11</sup> -GLP-1(7-37)OH	0.021
30 Val <sup>8</sup> -Lys <sup>11</sup> -GLP-1(7-37)OH	0.001
Val <sup>8</sup> -Tyr <sup>12</sup> -GLP-1(7-37)OH	0.81
Val <sup>8</sup> -Glu <sup>16</sup> -GLP-1(7-37)OH	0.047

- 86 -

	Val <sup>6</sup> -Ala <sup>16</sup> -GLP-1(7-37)OH	0.112
	Val <sup>8</sup> -Tyr <sup>16</sup> -GLP-1(7-37)OH	1.175
5	Val <sup>8</sup> -Lys <sup>20</sup> -GLP-1(7-37)OH	0.33
	Gln <sup>22</sup> -GLP-1(7-37)OH	0.42
10	Val <sup>6</sup> -Ala <sup>22</sup> -GLP-1(7-37)OH	0.56
	Val <sup>6</sup> -Ser <sup>22</sup> -GLP-1(7-37)OH	0.50
	Val <sup>6</sup> -Asp <sup>22</sup> -GLP-1(7-37)OH	0.40
15	Val <sup>6</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.29
	Val <sup>8</sup> -Lys <sup>22</sup> -GLP-1(7-37)OH	0.58
20	Val <sup>8</sup> -Pro <sup>22</sup> -GLP-1(7-37)OH	0.01
	Val <sup>8</sup> -His <sup>22</sup> -GLP-1(7-37)OH	0.14
	Val <sup>8</sup> -Lys <sup>22</sup> -GLP-1(7-36)NH <sub>2</sub>	0.53
25	Val <sup>6</sup> -Glu <sup>22</sup> -GLP-1(7-36)NH <sub>2</sub>	1.0
	Gly <sup>6</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.07
30	Val <sup>6</sup> -Lys <sup>23</sup> -GLP-1(7-37)OH	0.18
	Val <sup>6</sup> -His <sup>24</sup> -GLP-1(7-37)OH	0.007

- 87 -

	Val <sup>8</sup> -Lys <sup>24</sup> -GLP-1(7-37)OH	0.02
	Val <sup>8</sup> -His <sup>26</sup> -GLP-1(7-37)OH	1.6
5	Val <sup>8</sup> -Glu <sup>26</sup> -GLP-1(7-37)OH	1.5
	Val <sup>8</sup> -His <sup>27</sup> -GLP-1(7-37)OH	0.37
	Val <sup>8</sup> -Ala <sup>27</sup> -GLP-1(7-37)OH	0.47
10	Gly <sup>6</sup> -Glu <sup>30</sup> -GLP-1(7-37)OH	0.29
	Val <sup>8</sup> -Glu <sup>30</sup> -GLP-1(7-37)OH	0.29
15	Val <sup>8</sup> -Asp <sup>30</sup> -GLP-1(7-37)OH	0.15
	Val <sup>8</sup> -Ser <sup>30</sup> -GLP-1(7-37)OH	0.19
	Val <sup>8</sup> -His <sup>30</sup> -GLP-1(7-37)OH	0.19
20	Val <sup>8</sup> -Glu <sup>33</sup> -GLP-1(7-37)OH	0.039
	Val <sup>8</sup> -Ala <sup>33</sup> -GLP-1(7-37)OH	0.1
25	Val <sup>8</sup> -Gly <sup>33</sup> -GLP-1(7-37)OH	0.01
	Val <sup>8</sup> -Glu <sup>34</sup> -GLP-1(7-37)OH	0.17
	Val <sup>8</sup> -Pro <sup>35</sup> -GLP-1(7-37)OH	0.094
30	Val <sup>8</sup> -His <sup>35</sup> -GLP-1(7-37)OH	0.41
	Val <sup>8</sup> -Glu <sup>35</sup> -GLP-1(7-37)OH	0.15

- 88 -

	Val <sup>8</sup> -Glu <sup>36</sup> -GLP-1(7-37)OH	0.11
	Val <sup>8</sup> -His <sup>36</sup> -GLP-1(7-37)OH	0.22
5	Val <sup>8</sup> -His <sup>37</sup> -GLP-1(7-37)OH	0.33
	Val <sup>8</sup> -Leu <sup>16</sup> -Glu <sup>26</sup> -GLP-1(7-37)OH	0.23
10	Val <sup>8</sup> -Lys <sup>23</sup> -Glu <sup>30</sup> -GLP-1(7-37)OH	0.37
	Val <sup>8</sup> -Lys <sup>22</sup> -Glu <sup>23</sup> -GLP-1(7-37)OH	0.35
	Val <sup>8</sup> -Glu <sup>22</sup> -Ala <sup>27</sup> -GLP-1(7-37)OH	1.02
15	Val <sup>8</sup> -Glu <sup>22</sup> -Lys <sup>23</sup> -GLP-1(7-37)OH	1.43
	Val <sup>8</sup> -Lys <sup>33</sup> -Val <sup>34</sup> -GLP-1(7-37)OH	0.08
20	Val <sup>8</sup> -Lys <sup>33</sup> -Asn <sup>34</sup> -GLP-1(7-37)OH	0.09
	Val <sup>8</sup> -Gly <sup>34</sup> -Lys <sup>35</sup> -GLP-1(7-37)OH	0.34
25	Val <sup>8</sup> -Gly <sup>36</sup> -Pro <sup>37</sup> -GLP-1(7-37)NH <sub>2</sub>	0.53

Table 5b. *In vitro* GLP-1 analog activity

GLP-1 Compound	GLP-1 Receptor Activation
GLP-1(7-37)OH	1.0
Val <sup>8</sup> -GLP-1(7-37)OH	0.47
Gly <sup>8</sup> -GLP-1(7-37)OH	0.80
Val <sup>8</sup> -Tyr <sup>12</sup> -GLP-1(7-37)OH	0.80
Val <sup>8</sup> -Tyr <sup>12</sup> -GLP-1(7-36)NH <sub>2</sub>	0.52
Val <sup>8</sup> -Trp <sup>12</sup> -GLP-1(7-37)OH	0.52
Val <sup>8</sup> -Leu <sup>16</sup> -GLP-1(7-37)OH	0.52
Val <sup>8</sup> -Val <sup>16</sup> -GLP-1(7-37)OH	0.52
Val <sup>8</sup> -Tyr <sup>16</sup> -GLP-1(7-37)OH	1.18
Gly <sup>8</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.03
Val <sup>8</sup> -Leu <sup>25</sup> -GLP-1(7-37)OH	0.24
Val <sup>8</sup> -Tyr <sup>12</sup> -Tyr <sup>16</sup> -GLP-1(7-37)OH	0.70
Val <sup>8</sup> -Trp <sup>12</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	0.80
Val <sup>8</sup> -Tyr <sup>12</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.27
Val <sup>8</sup> -Tyr <sup>16</sup> -Phe <sup>19</sup> -GLP-1(7-37)OH	1.32
Val <sup>8</sup> -Tyr <sup>16</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.69, 1.79
Val <sup>8</sup> -Trp <sup>16</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	2.30, 2.16
Val <sup>8</sup> -Leu <sup>16</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	2.02
Val <sup>8</sup> -Ile <sup>16</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.55
Val <sup>8</sup> -Phe <sup>16</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.08
Val <sup>8</sup> -Trp <sup>19</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.50, 3.10
Val <sup>8</sup> -Tyr <sup>18</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	2.40, 2.77
Val <sup>8</sup> -Phe <sup>18</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	0.94
Val <sup>8</sup> -Ile <sup>18</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.88
Val <sup>8</sup> -Lys <sup>18</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.18
Val <sup>8</sup> -Trp <sup>19</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.50
Val <sup>8</sup> -Phe <sup>19</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	0.70
Val <sup>8</sup> -Phe <sup>20</sup> -Glu <sup>22</sup> -GLP-1(7-37)OH	1.27

- 90 -

Val <sup>8</sup> -Glu <sup>22</sup> -Leu <sup>25</sup> -GLP-1(7-37)OH	1.32
Val <sup>8</sup> -Glu <sup>22</sup> -Ile <sup>45</sup> -GLP-1(7-37)OH	1.46
Val <sup>8</sup> -Glu <sup>22</sup> -Val <sup>25</sup> -GLP-1(7-37)OH	2.21, 1.36
Val <sup>8</sup> -Glu <sup>22</sup> -Ile <sup>27</sup> -GLP-1(7-37)OH	0.94
Val <sup>8</sup> -Glu <sup>22</sup> -Ala <sup>27</sup> -GLP-1(7-37)OH	1.03
Val <sup>8</sup> -Glu <sup>22</sup> -Ile <sup>33</sup> -GLP-1(7-37)OH	2.21, 1.79, 1.60
Val <sup>8</sup> -Asp <sup>9</sup> -Ile <sup>11</sup> -Tyr <sup>16</sup> -Glu <sup>22</sup> - GLP-1(7-37)OH	2.02
Val <sup>8</sup> -Tyr <sup>16</sup> -Trp <sup>19</sup> -Glu <sup>22</sup> -GLP-1(7- 37)OH	1.64
Val <sup>8</sup> -Trp <sup>16</sup> -Glu <sup>22</sup> -Val <sup>25</sup> -Ile <sup>33</sup> - GLP-1(7-37)OH	2.35
Val <sup>8</sup> -Trp <sup>16</sup> -Glu <sup>22</sup> -Ile <sup>33</sup> -GLP-1(7- 37)OH	1.93
Val <sup>8</sup> -Glu <sup>22</sup> -Val <sup>25</sup> -Ile <sup>33</sup> -GLP-1(7- 37)OH	2.30, 2.73, 3.15
Val <sup>8</sup> -Trp <sup>16</sup> -Glu <sup>22</sup> -Val <sup>25</sup> -GLP-1(7- 37)OH	2.07
Val <sup>8</sup> -Cys <sup>16</sup> -Lys <sup>45</sup> -GLP-1(7-37)OH	1.97
Val <sup>8</sup> -Cys <sup>16</sup> -Lys <sup>46</sup> -Arg <sup>38</sup> -GLP-1(7- 37)OH	2.4, 1.9

Table 6: In vitro activity of GLP/Exendin analogs

Peptide Sequence	In Vitro Activity (% of Val <sup>8</sup> -GLP- 1(7-37)OH)
HGEGTFTSDLSKQMEEEAVRLFIEWLKNGGP-NH <sub>2</sub>	6.21
HGEGTFTSDLSKQMEEEAVRLFIEWLKNGGPSSGAPP PS-NH <sub>2</sub>	6.75, 3.25
HVEGTFTSDLSKQMEEEAVRLFIAWLKGRG	2.86
HVEGTFTSDVSSYLEEEAVRLFIAWLKGRG	1.47
HVEGTFTSDLSKQMEGQAQKEFIAWLKGRG	0.11

- 91 -

HVEGTFTSDVSKQMEGQAAKEFIAWLVKGRG	0.04
HGEGTFTSDLSKQMEGQAAKEFIEWLKNGGP-NH2	1.44
HGEGTFTSDLSKQMEEEAAKEFIEWLKNGGP-NH2	2.80
HGEGTFTSDVSSYLEEEAVRLFIEWLKNGGP-NH2	5.40
HGEGTFTSDLSYLEEEAVRLFIEWLKNGGP-NH2	5.07
HAEGTFTSDVSSYLEGQAAKEFIAWLVKGRPSSGAPPPS-NH2	3.30
HAEGTFTSDVSKQLEEEAAKEFIAWLVKGRG	2.15
HVEGTFTSDVSSYLEGQAAKEFIEWLKNGGP-NH2	2.36
HGEGTFTSDLSKQMEEEAVRLFIAWLVKGRG	3.25
HVEGTFTSDVSSYLEEEAAKEFIAWLVKGRG	1.00
HVEGTFTSDVSSYLEGQAAKEFIAWLKNGRG	0.20
HVEGTFTSDVSSYLEGQAAKEFIAWLVKGRG	1.00
HAEGTFTSDVSSYLEGQAAKEFIAWLVKGRG	2.12

Example 7: *In vivo* pharmacokinetics of Val<sup>8</sup>-GLP-1-IgG1  
and Val<sup>8</sup>-GLP-1-HSA:

- A pharmacokinetic study of Val<sup>8</sup>-GLP-1-IgG1 and Val<sup>8</sup>-GLP-1-HSA was performed in cynomolgus monkeys. Monkeys were dosed at 5.6 nmoles/kg with either purified Val<sup>8</sup>-GLP-1-IgG1 or Val<sup>8</sup>-GLP-1-HSA. The compounds were administered as an intravenous bolus administration. Blood was collected pre-dose and at 0.083, 0.25, 0.5, 1, 4, 8, 12, 24, 48, 72, 96, 120, 144, 168, and 216 hours post-dose into tubes containing EDTA. Plasma concentrations of immunoreactive Val<sup>8</sup>-GLP-1 were determined using a radioimmunoassay that utilizes a polyclonal antiserum whose primary specificity is for the N-terminal (7-16) region of Val<sup>8</sup>-GLP-1(7-37).
- Figure 9 depicts the plasma concentration of Val<sup>8</sup>-GLP-1-Fc and Val<sup>8</sup>-GLP-1-Linker-HSA following a single intravenous dose to two cynomolgus monkeys. The Fc fusion protein had a half-life of approximately 45 hours and the albumin fusion had a half-life of approximately 87 hours.



- 92 -

Example 8: In vivo pharmacodynamics of Exendin-4-IgG1:

Two chronically cannulated normal male beagle dogs were studied after an overnight fast. Arterial and venous vascular access ports were accessed, and a catheter was inserted percutaneously into a cephalic vein and secured. Animals were placed in cages, and their catheters were attached to a swivel/tether system. A solution containing the fusion protein Exendin-4-IgG1 (11.8  $\mu$ M) was injected intravenously (1.0 nmol/kg) through the cephalic vein catheter. The catheter was then cleared with 10 ml of saline. Two hours later, a hyperglycemic (150 mg/dl) clamp was initiated and continued for three hours. Arterial blood samples were drawn throughout this 5-hour period for determination of plasma concentrations of the fusion protein, glucose, and insulin.

The results of this study were compared to those from a similar, previous study in which both of the animals had received a bolus of saline, s.c., and three hours later were studied using a 3-hour hyperglycemic (150 mg/dl) clamp.

In both sets of studies, plasma glucose concentrations were determined using a Beckman glucose analyzer. Plasma insulin concentrations were determined by employees of Linco Research, Inc. using an RIA kit developed in their laboratories. The data is illustrated in Figures 10 and 11.

Example 9: In vivo pharmacokinetics ofGly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-IgG1:

Two groups of three normal male beagle dogs received 0.1 mg/kg of Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-IgG1 by subcutaneous (SC) or intravenous (IV) administration. Plasma concentrations of Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-IgG1 immunoreactivity were determined by radioimmunoassay in samples collected from 30 minutes predose to 216 hours postdose for both the IV and SC groups. These concentrations were subsequently used to determine the

- 93 -

reported pharmacokinetic parameters. The mean elimination half-life of IV administered Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-IgG1 was approximately 55 hours and the total body clearance was 1.5 mL/h/kg. The mean elimination half-life of SC  
5 administered Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1-CEX-Linker-IgG1 was approximately 38 hours.

- 94 -

WE CLAIM:

1. A heterologous fusion protein comprising a first  
5 polypeptide with a N-terminus and a C-terminus fused to a  
second polypeptide with a N-terminus and a C-terminus  
wherein the first polypeptide is a GLP-1 compound and the  
second polypeptide is selected from the group consisting of
  - a) human albumin;
  - 10 b) human albumin analogs; and
  - c) fragments of human albumin,and wherein the C-terminus of the first polypeptide is fused  
to the N-terminus of the second polypeptide.
- 15 2. A heterologous fusion protein comprising a first  
polypeptide with a N-terminus and a C-terminus fused to a  
second polypeptide with a N-terminus and a C-terminus  
wherein the first polypeptide is a GLP-1 compound and the  
second polypeptide is selected from the group consisting of
  - 20 a) human albumin;
  - b) human albumin analogs; and
  - c) fragments of human albumin,and wherein the C-terminus of the first polypeptide is fused  
to the N-terminus of the second polypeptide via a peptide  
25 linker.
3. The heterologous fusion protein of the Claim 2 wherein  
the peptide linker is selected from the group consisting of:
  - a) a glycine rich peptide;
  - 30 b) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>n</sub>,  
where n is 1, 2, 3, 4, 5 or 6; and
  - c) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>3</sub>.

- 95 -

4. The heterologous fusion protein of Claims 1, 2, or 3 wherein the GLP-1 compound comprises the sequence of formula 1 [SEQ ID NO: 2]

5           7   8   9   10   11   12   13   14   15   16   17  
          His-Xaa-Xaa-Gly-Xaa-Phe-Thr-Xaa-Asp-Xaa-Xaa-  
          18 19 20 21 22 23 24 25 26 27 28  
          Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Phe-  
          29 30 31 32 33 34 35 36 37 38 39  
10       Ile-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-  
          40 41 42 43 44 45  
          Xaa-Xaa-Xaa-Xaa-Xaa-Xaa

Formula I (SEQ ID NO: 2)

15           wherein:

Xaa at position 8 is Ala, Gly, Ser, Thr, Leu, Ile, Val,  
          Glu, Asp, or Lys;  
Xaa at position 9 is Glu, Asp, or Lys;  
Xaa at position 11 is Thr, Ala, Gly, Ser, Leu, Ile,  
20       Val, Glu, Asp, or Lys;  
Xaa at position 14 is Ser, Ala, Gly, Thr, Leu, Ile,  
          Val, Glu, Asp, or Lys;  
Xaa at position 16 is Val, Ala, Gly, Ser, Thr, Leu,  
          Ile, Tyr, Glu, Asp, Trp, or Lys;  
25       Xaa at position 17 is Ser, Ala, Gly, Thr, Leu, Ile,  
          Val, Glu, Asp, or Lys;  
Xaa at position 18 is Ser, Ala, Gly, Thr, Leu, Ile,  
          Val, Glu, Asp, Trp, Tyr, or Lys;  
Xaa at position 19 is Tyr, Phe, Trp, Glu, Asp, Gln, or  
30       Lys;  
Xaa at position 20 is Leu, Ala, Gly, Ser, Thr, Ile,  
          Val, Glu, Asp, Met, Trp, Tyr, or Lys;  
Xaa at position 21 is Glu, Asp, or Lys;  
Xaa at position 22 is Gly, Ala, Ser, Thr, Leu, Ile,  
35       Val, Glu, Asp, or Lys;

- 96 -

- Xaa at position 23 is Gln, Asn, Arg, Glu, Asp, or Lys;  
Xaa at position 24 is Ala, Gly, Ser, Thr, Leu, Ile,  
Val, Arg, Glu, Asp, or Lys;  
Xaa at position 25 is Ala, Gly, Ser, Thr, Leu, Ile,  
5 Val, Glu, Asp, or Lys;  
Xaa at position 26 is Lys, Arg, Gln, Glu, Asp, or His;  
Xaa at position 27 is Leu, Glu, Asp, or Lys;  
Xaa at position 30 is Ala, Gly, Ser, Thr, Leu, Ile,  
Val, Glu, Asp, or Lys;  
10 Xaa at position 31 is Trp, Phe, Tyr, Glu, Asp, or Lys;  
Xaa at position 32 is Leu, Gly, Ala, Ser, Thr, Ile,  
Val, Glu, Asp, or Lys;  
Xaa at position 33 is Val, Gly, Ala, Ser, Thr, Leu,  
Ile, Glu, Asp, or Lys;  
15 Xaa at position 34 is Asn, Lys, Arg, Glu, Asp, or His;  
Xaa at position 35 is Gly, Ala, Ser, Thr, Leu, Ile,  
Val, Glu, Asp, or Lys;  
Xaa at position 36 is Gly, Arg, Lys, Glu, Asp, or His;  
Xaa at position 37 is Pro, Gly, Ala, Ser, Thr, Leu,  
20 Ile, Val, Glu, Asp, or Lys, or is deleted;  
Xaa at position 38 is Ser, Arg, Lys, Glu, Asp, or His,  
or is deleted;  
Xaa at position 39 is Ser, Arg, Lys, Glu, Asp, or His,  
or is deleted;  
25 Xaa at position 40 is Gly, Asp, Glu, or Lys, or is  
deleted;  
Xaa at position 41 is Ala, Phe, Trp, Tyr, Glu, Asp, or  
Lys, or is deleted;  
Xaa at position 42 is Ser, Pro, Lys, Glu, or Asp, or is  
30 deleted;  
Xaa at position 43 is Ser, Pro, Glu, Asp, or Lys, or is  
deleted;  
Xaa at position 44 is Gly, Pro, Glu, Asp, or Lys, or is  
deleted; and

- 97 -

Xaa at position 45 is Ala, Ser, Val, Glu, Asp, or Lys,  
or is

deleted;

provided that when the amino acid at position 37, 38, 39,  
5 40, 41, 42, 43, or 44 is deleted, then each amino acid  
downstream of that amino acid is also deleted.

5. The heterologous fusion protein of Claims 1, 2, or 3  
wherein the GLP-1 compound comprises the sequence of formula  
10 II (SEQ ID NO: 3):

7 8 9 10 11 12 13 14 15 16 17  
Xaa-Xaa-Xaa-Gly-Xaa-Xaa-Thr-Ser-Asp-Xaa-Ser-  
18 19 20 21 22 23 24 25 26 27 28  
15 Xaa-Tyr-Leu-Glu-Xaa-Xaa-Xaa-Ala-Xaa-Xaa-Phe-  
29 30 31 32 33 34 35 36 37  
Ile-Xaa-Xaa-Leu-Xaa-Xaa-Xaa-Xaa-Xaa  
Formula II (SEQ ID NO: 3)

20 wherein:

Xaa at position 7 is: L-histidine, D-histidine,  
desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-  
histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  
 $\alpha$ -methyl-histidine;

25 Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or  
Thr;

Xaa at position 9 is: Thr, Ser, Arg, Lys, Trp, Phe,  
Tyr, Glu, or His;

Xaa at position 11 is: Asp, Glu, Arg, Thr, Ala, Lys, or His;

30 Xaa at position 12 is: His, Trp, Phe, or Tyr;

Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp,  
Val, Tyr, Glu, or Ala;

Xaa at position 18 is: His, Pro, Asp, Glu, Arg, Ser, Ala, or  
Lys;

35 Xaa at position 19 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or

- 98 -

Cys;

Xaa at position 23 is: His, Asp, Lys, Glu, Gln, or Arg;

Xaa at position 24 is: Glu, Arg, Ala, or Lys;

Xaa at position 26 is: Trp, Tyr, Phe, Asp, Lys, Glu, or His;

5 Xaa at position 27 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys;

Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;

Xaa at position 31 is: Asp, Glu, Ser, Thr, Arg, Trp, or Lys;

Xaa at position 33 is: Asp, Arg, Val, Lys, Ala, Gly, or Glu;

10 Xaa at position 34 is: Glu, Lys, or Asp;

Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;

Xaa at position 36 is: Thr, Ser, Asp, Trp, Tyr, Phe, Arg, Glu, or His;

15 Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

6. The heterologous fusion protein of Claims 1, 2, or 3 wherein the GLP-1 compound comprises the sequence of formula

20 III (SEQ ID NO: 4):

```

      7   8   9   10  11  12  13  14  15  16  17
Xaa-Xaa-Glu-Gly-Xaa-Xaa-Thr-Ser-Asp-Xaa-Ser-
18 19 20 21 22 23 24 25 26 27 28
Ser-Tyr-Leu-Glu-Xaa-Xaa-Xaa-Ala-Xaa-Xaa-Phe-
25 29 30 31 32 33 34 35 36 37
Ile-Ala-Xaa-Leu-Xaa-Xaa-Xaa-Xaa

```

formula III (SEQ ID NO: 4)

wherein:

30 Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;

Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;

35 Xaa at position 11 is: Asp, Glu, Arg, Thr, Ala, Lys, or His;

- 99 -

Xaa at position 12 is: His, Trp, Phe, or Tyr;

Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp,  
Val, Glu, or Ala;

Xaa at position 22: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or  
5 Cys;

Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;

Xaa at position 24 is: Glu, His, Ala, or Lys;

Xaa at position 25 is: Asp, Lys, Glu, or His;

Xaa at position 27 is: Ala, Glu, His, Phe, Tyr, Trp, Arg,  
10 or Lys;

Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;

Xaa at position 33 is: Asp, Arg, Val, Lys, Ala, Gly, or  
Glu;

Xaa at position 34 is: Glu, Lys, or Asp;

15 Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe,  
Asp, Gly, Pro, His, or Glu;

Xaa at position 36 is: Arg, Glu, or His;

Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp,  
Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

20

7. The heterologous fusion protein of Claim 1, 2, or 3  
wherein the GLP-1 compound comprises the sequence of formula  
IV (SEQ ID NO: 5):

	7	8	9	10	11	12	13	14	15	16	17
25	Xaa-Xaa-Glu-Gly-Thr-Xaa-Thr-Ser-Asp-Xaa-Ser-										
	18	19	20	21	22	23	24	25	26	27	28
	Ser-Tyr-Leu-Glu-Xaa-Xaa-Ala-Ala-Xaa-Glu-Phe-										
	29	30	31	32	33	34	35	36	37		
	Ile-Xaa-Trp-Leu-Val-Lys-Xaa-Arg-Xaa										
30	formula IV (SEQ ID NO: 5)										

wherein:

Xaa at position 7 is: L-histidine, D-histidine, desamino-  
histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine,



- 100 -

homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;

Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;

Xaa at position 12 is: His, Trp, Phe, or Tyr;

5 Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp, Val, Glu, or Ala;

Xaa at position 22 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;

Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;

10 Xaa at position 26 is: Asp, Lys, Glu, or His;

Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;

Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;

15 Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His,  $-\text{NH}_2$ , Gly, Gly-Pro, or Gly-Pro- $\text{NH}_2$ , or is deleted.

8. The heterologous fusion protein of Claims 1, 2, or 3 wherein the GLP-1 compound comprises the sequence of formula

20 V (SEQ ID NO: 6)

7 8 9 10 11 12 13 14 15 16 17  
Xaa-Xaa-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-  
18 19 20 21 22 23 24 25 26 27 28  
25 Ser-Tyr-Leu-Glu-Xaa-Xaa-Ala-Ala-Lys-Xaa-Phe-  
29 30 31 32 33 34 35 36 37  
Ile-Xaa-Trp-Leu-Val-Lys-Gly-Arg-Xaa  
formula V (SEQ ID NO: 6)

30 wherein:

Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;

35 Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;

- 101 -

Xaa at position 22 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;

Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;

Xaa at position 24 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys

Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;

Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

- 10 9. The heterologous fusion protein of Claims 1, 2, or 3 wherein the GLP-1 compound comprises the sequence of formula VIII (SEQ ID NO:11).

15                   7   8   9   10 11 12 13 14 15 16 17  
Xaa-Xaa-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Xaa-Ser-  
18 19 20 21 22 23 24 25 26 27 28  
Xaa-Xaa-Xaa-Glu-Xaa-Xaa-Ala-Xaa-Xaa-Xaa-Phe-  
29 30 31 32 33 34 35 36 37  
Ile-Xaa-Trp-Leu-Xaa-Xaa-Gly-Xaa-Xaa  
20                   formula VIII (SEQ ID NO: 11)

wherein:

- Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine,  
25       homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;  
Xaa at position 8 is: Gly, Ala, or Val;  
Xaa at position 16 is: Leu or Val;  
Xaa at position 18 is Lys or Ser;  
30 Xaa at position 19 is: Gln or Tyr;  
Xaa at position 20 is: Met or Leu;  
Xaa at position 22 is: Glu or Gln;  
Xaa at position 23 is: Glu, or Gln;  
Xaa at position 25 is: Val or Ala;  
35 Xaa at position 26 is: Arg or Lys;

- 102 -

- Xaa at position 27 is Leu or Glu;  
Xaa at position 30 is: Glu or Ala;  
Xaa at position 33 is: Val or Lys;  
Xaa at position 34 is: Asn or Lys;  
5 Xaa at position 36 is: Gly or Arg; and  
Xaa at position 37 is: Gly, Pro, Pro-Ser-Ser-Gly-Ala-Pro-  
Pro-Pro-Ser, or is absent.
10. The heterologous fusion protein of claims 1 through 9  
10 wherein the GLP-1 compound has no more than 6 amino acids  
that are different from the corresponding amino acid in GLP-  
1(7-37)OH, GLP-1(7-36)OH, or Exendin-4.
11. The heterologous fusion protein of Claim 10 wherein the  
15 GLP-1 compound has no more than 5 amino acids that differ  
from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-  
36)OH, or Exendin-4.
12. The heterologous fusion protein of Claim 11 wherein the  
20 GLP-1 compound has no more than 4 amino acids that differ  
from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-  
36)OH, or Exendin-4.
13. The heterologous fusion protein of Claim 12 wherein the  
25 GLP-1 compound has no more than 3 amino acids that differ  
from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-  
36)OH, or Exendin-4.
14. The heterologous fusion protein of Claim 13 wherein the  
30 GLP-1 compound has no more than 2 amino acids that differ  
from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-  
36)OH, or Exendin-4.
15. The heterologous fusion protein of any one of Claims 1  
35 through 14 wherein Xaa at position 8 is glycine or valine.

- 103 -

16. The heterologous fusion protein of Claim 8 wherein the GLP-1 compound has no more than 2 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH or GLP-1(7-36)OH and Xaa at position 8 is glycine or valine and Xaa at position 30 is alanine, glutamic acid, aspartic acid, serine, or histidine.
17. The heterologous fusion protein of Claim 16 wherein Xaa at position 30 is Glutamic acid.
18. The heterologous fusion protein of Claim 8 wherein the GLP-1 compound has no more than 2 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH or GLP-1(7-36)OH and Xaa at position 8 is glycine or valine and Xaa at position 37 is histidine, phenylalanine, tyrosine, tryptophan, aspartic acid, glutamic acid, serine, threonine, arginine, or lysine.
19. The heterologous fusion protein of Claim 18 wherein Xaa at position 37 is histidine.
20. The heterologous fusion protein of Claim 8 wherein the GLP-1 compound has no more than 2 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH or GLP-1(7-36)OH and Xaa at position 8 is glycine, valine, leucine, isoleucine, serine, threonine, or methionine and Xaa at position 22 is aspartic acid, glutamic acid, lysine, arginine, asparagine, glutamine or histidine.
21. The heterologous fusion protein of Claim 20 wherein Xaa at position 22 is lysine or glutamic acid.
22. The heterologous fusion protein of Claim 8 wherein the GLP-1 compound is Val<sup>8</sup>-GLP-1(7-37).

- 104 -

23. The heterologous fusion protein of Claim 8 wherein the GLP-1 compound is Gly<sup>B</sup>-GLP-1(7-37).

5 24. The heterologous fusion protein of any one of Claims 1 through 23 wherein the second polypeptide is human albumin.

25. The heterologous fusion protein of Claim 24 wherein the second polypeptide has the sequence of SEQ ID NO: 34.

10

26. The heterologous fusion protein of any one of Claims 1 through 23 wherein the second polypeptide is an N-terminal fragment of albumin.

15 27. A heterologous fusion protein comprising a first polypeptide with a N-terminus and a C-terminus fused to a second polypeptide with a N-terminus and a C-terminus wherein the first polypeptide is a GLP-1 compound and the second polypeptide is selected from the group consisting of  
20 a) the Fc portion of an immunoglobulin;  
b) an analog of the Fc portion of an immunoglobulin;  
and  
c) fragments of the Fc portion of an immunoglobulin,  
and wherein the C-terminus of the first polypeptide is fused  
25 to the N-terminus of the second polypeptide.

28. A heterologous fusion protein comprising a first polypeptide with a N-terminus and a C-terminus fused to a second polypeptide with a N-terminus and a C-terminus  
30 wherein the first polypeptide is a GLP-1 compound and the second polypeptide is selected from the group consisting of  
a) the Fc portion of an immunoglobulin;  
b) an analog of the Fc portion of an immunoglobulin;  
and  
35

- 105 -

c) fragments of the Fc portion of an immunoglobulin,  
and wherein the C-terminus of the first polypeptide is  
fused to the N-terminus of the second polypeptide via a  
5 peptide linker.

29. The heterologous fusion protein of the Claim 28 wherein  
the peptide linker is selected from the group consisting of:  
a) a glycine rich peptide;  
10 b) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>n</sub>  
where n is 1, 2, 3, 4, 5, or 6; and  
c) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]<sub>3</sub>.
30. The heterologous fusion protein of any one of Claims  
15 27, 28, or 29 wherein the GLP-1 compound comprises the  
sequence of formula 1 [SEQ ID NO: 2].

7 8 9 10 11 12 13 14 15 16 17  
His-Xaa-Xaa-Gly-Xaa-Phe-Thr-Xaa-Asp-Xaa-Xaa-  
20 18 19 20 21 22 23 24 25 26 27 28  
Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Phe-  
29 30 31 32 33 34 35 36 37 38 39  
Ile-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-  
40 41 42 43 44 45  
25 Xaa-Xaa-Xaa-Xaa-Xaa-Xaa

Formula I (SEQ ID NO: 2)

wherein:

- Xaa at position 8 is Ala, Gly, Ser, Thr, Leu, Ile, Val,  
30 Glu, Asp, or Lys;  
Xaa at position 9 is Glu, Asp, or Lys;  
Xaa at position 11 is Thr, Ala, Gly, Ser, Leu, Ile, Val,  
Glu, Asp, or Lys;  
Xaa at position 14 is Ser, Ala, Gly, Thr, Leu, Ile, Val,  
35 Glu, Asp, or Lys;

- 106 -

- Xaa at position 16 is Val, Ala, Gly, Ser, Thr, Leu, Ile, Tyr, Glu, Asp, Trp, or Lys;
- Xaa at position 17 is Ser, Ala, Gly, Thr, Leu, Ile, Val, Glu, Asp, or Lys;
- 5 Xaa at position 18 is Ser, Ala, Gly, Thr, Leu, Ile, Val, Glu, Asp, Trp, Tyr, or Lys;
- Xaa at position 19 is Tyr, Phe, Trp, Glu, Asp, Gln, or Lys;
- Xaa at position 20 is Leu, Ala, Gly, Ser, Thr, Ile, Val, Glu, Asp, Met, Trp, Tyr, or Lys;
- 10 Xaa at position 21 is Glu, Asp, or Lys;
- Xaa at position 22 is Gly, Ala, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;
- Xaa at position 23 is Gln, Asn, Arg, Glu, Asp, or Lys;
- Xaa at position 24 is Ala, Gly, Ser, Thr, Leu, Ile, Val,
- 15 Arg, Glu, Asp, or Lys;
- Xaa at position 25 is Ala, Gly, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;
- Xaa at position 26 is Lys, Arg, Gln, Glu, Asp, or His;
- Xaa at position 27 is Leu, Glu, Asp, or Lys;
- 20 Xaa at position 30 is Ala, Gly, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;
- Xaa at position 31 is Trp, Phe, Tyr, Glu, Asp, or Lys;
- Xaa at position 32 is Leu, Gly, Ala, Ser, Thr, Ile, Val, Glu, Asp, or Lys;
- 25 Xaa at position 33 is Val, Gly, Ala, Ser, Thr, Leu, Ile, Glu, Asp, or Lys;
- Xaa at position 34 is Asn, Lys, Arg, Glu, Asp, or His;
- Xaa at position 35 is Gly, Ala, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;
- 30 Xaa at position 36 is Gly, Arg, Lys, Glu, Asp, or His;
- Xaa at position 37 is Pro, Gly, Ala, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys, or is deleted;
- Xaa at position 38 is Ser, Arg, Lys, Glu, Asp, or His, or is deleted;
- 35 Xaa at position 39 is Ser, Arg, Lys, Glu, Asp, or His, or

- 107 -

is deleted;

Xaa at position 40 is Gly, Asp, Glu, or Lys, or is deleted;

Xaa at position 41 is Ala, Phe, Trp, Tyr, Glu, Asp, or Lys,  
or is deleted;

5 Xaa at position 42 is Ser, Pro, Lys, Glu, or Asp, or is  
deleted;

Xaa at position 43 is Ser, Pro, Glu, Asp, or Lys, or is  
deleted;

10 Xaa at position 44 is Gly, Pro, Glu, Asp, or Lys, or is  
deleted;

and

Xaa at position 45 is Ala, Ser, Val, Glu, Asp, or Lys, or is  
deleted;

15 provided that when the amino acid at position 37, 38, 39,  
40, 41, 42, 43, or 44 is deleted, then each amino acid  
downstream of that amino acid is also deleted.

31. The heterologous fusion protein of Claims 27, 28, or 29  
wherein the GLP-1 compound comprises the sequence of formula  
20 II (SEQ ID NO: 3):

7 8 9 10 11 12 13 14 15 16 17  
Xaa-Xaa-Xaa-Gly-Xaa-Xaa-Thr-Ser-Asp-Xaa-Ser-  
18 19 20 21 22 23 24 25 26 27 28  
25 Xaa-Tyr-Leu-Glu-Xaa-Xaa-Xaa-Ala-Xaa-Xaa-Phe-  
29 30 31 32 33 34 35 36 37  
Ile-Xaa-Xaa-Leu-Xaa-Xaa-Xaa-Xaa-Xaa

Formula II (SEQ ID NO: 3)

30 wherein:

Xaa at position 7 is: L-histidine, D-histidine, desamino-  
histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine,  
homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-  
histidine;

35 Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;



- 108 -

- Xaa at position 9 is: Thr, Ser, Arg, Lys, Trp, Phe, Tyr,  
Glu, or His;
- Xaa at position 11 is: Asp, Glu, Arg, Thr, Ala, Lys, or His;
- Xaa at position 12 is: His, Trp, Phe, or Tyr;
- 5 Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp,  
Val, Tyr, Glu, or Ala;
- Xaa at position 18 is: His, Pro, Asp, Glu, Arg, Ser, Ala, or  
Lys;
- Xaa at position 19 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or  
10 Cys;
- Xaa at position 23 is: His, Asp, Lys, Glu, Gln, or Arg;
- Xaa at position 24 is: Glu, Arg, Ala, or Lys;
- Xaa at position 26 is: Trp, Tyr, Phe, Asp, Lys, Glu, or His;
- Xaa at position 27 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or  
15 Lys;
- Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;
- Xaa at position 31 is: Asp, Glu, Ser, Thr, Arg, Trp, or Lys;
- Xaa at position 33 is: Asp, Arg, Val, Lys, Ala, Gly, or Glu;
- Xaa at position 34 is: Glu, Lys, or Asp;
- 20 Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe,  
Asp, Gly, Pro, His, or Glu;
- Xaa at position 36 is: Thr, Ser, Asp, Trp, Tyr, Phe, Arg,  
Glu, or His;
- Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp,  
25 Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

32. The heterologous fusion protein of Claims 27, 28, or 29  
wherein the GLP-1 compound comprises the sequence of formula  
III (SEQ ID NO: 4):

- |    |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 30 | 7   | 8   | 9   | 10  | 11  | 12  | 13  | 14  | 15  | 16  | 17  |
|    | Xaa | Xaa | Glu | Gly | Xaa | Xaa | Thr | Ser | Asp | Xaa | Ser |
|    | 18  | 19  | 20  | 21  | 22  | 23  | 24  | 25  | 26  | 27  | 28  |
|    | Ser | Tyr | Leu | Glu | Xaa | Xaa | Xaa | Ala | Xaa | Xaa | Phe |
|    | 29  | 30  | 31  | 32  | 33  | 34  | 35  | 36  | 37  |     |     |
| 35 | Ile | Ala | Xaa | Leu | Xaa | Xaa | Xaa | Xaa | Xaa | R   |     |

- 109 -

## Formula III (SEQ ID NO: 4)

wherein:

- 5 Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;
- Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;
- Xaa at position 11 is: Asp, Glu, Arg, Thr, Ala, Lys, or His;
- 10 Xaa at position 12 is: His, Trp, Phe, or Tyr;
- Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp, Val, Glu, or Ala;
- Xaa at position 22: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;
- 15 Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;
- Xaa at position 24 is: Glu, His, Ala, or Lys;
- Xaa at position 25 is: Asp, Lys, Glu, or His;
- Xaa at position 27 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys;
- 20 Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;
- Xaa at position 33 is: Asp, Arg, Val, Lys, Ala, Gly, or Glu;
- Xaa at position 34 is: Glu, Lys, or Asp;
- Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;
- 25 Xaa at position 36 is: Arg, Glu, or His;
- Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.
- 30 33. The heterologous fusion protein of Claim 27, 28, or 29 wherein the GLP-1 compound comprises the sequence of formula IV (SEQ ID NO: 5):
- |    |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
|    | 7   | 8   | 9   | 10  | 11  | 12  | 13  | 14  | 15  | 16  | 17  |
|    | Xaa | Xaa | Glu | Gly | Thr | Xaa | Thr | Ser | Asp | Xaa | Ser |
| 35 | 18  | 19  | 20  | 21  | 22  | 23  | 24  | 25  | 26  | 27  | 28  |

- 110 -

Ser-Tyr-Leu-Glu-Xaa-Xaa-Ala-Ala-Xaa-Glu-Phe-  
29 30 31 32 33 34 35 36 37  
Ile-Xaa-Trp-Leu-Val-Lys-Xaa-Arg-Xaa  
formula IV (SEQ ID NO: 5)

5

wherein:

- Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;  
10 Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;  
Xaa at position 12 is: His, Trp, Phe, or Tyr;  
Xaa at position 16 is: Leu, Ser, Thr, Trp, His, Phe, Asp, Val, Glu, or Ala;  
15 Xaa at position 22 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;  
Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;  
Xaa at position 26 is: Asp, Lys, Glu, or His;  
Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;  
20 Xaa at position 35 is: Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;  
Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

- 25 34. The heterologous fusion protein of Claims 27, 28, or 29 wherein the GLP-1 compound comprises the sequence of formula V (SEQ ID NO: 6)

- 7 8 9 10 11 12 13 14 15 16 17  
30 Xaa-Xaa-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-  
18 19 20 21 22 23 24 25 26 27 28  
Ser-Tyr-Leu-Glu-Xaa-Xaa-Ala-Ala-Lys-Xaa-Phe-  
29 30 31 32 33 34 35 36 37  
Ile-Xaa-Trp-Leu-Val-Lys-Gly-Arg-Xaa  
35 formula V (SEQ ID NO: 6)

- 111 -

wherein:

- Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;
- 5 Xaa at position 8 is: Gly, Ala, Val, Leu, Ile, Ser, or Thr;
- Xaa at position 22 is: Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;
- 10 Xaa at position 23 is: His, Asp, Lys, Glu, or Gln;
- Xaa at position 24 is: Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys
- Xaa at position 30 is: Ala, Glu, Asp, Ser, or His;
- Xaa at position 37 is: Lys, Arg, Thr, Ser, Glu, Asp, Trp,
- 15 Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

35. The heterologous fusion protein of Claims 27, 28, or 29 wherein the GLP-1 compound comprises the sequence of formula VIII (SEQ ID NO:11).

- 20
- |    |     |     |     |     |     |     |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
|    | 7   | 8   | 9   | 10  | 11  | 12  | 13  | 14  | 15  | 16  | 17  |
|    | Xaa | Xaa | Glu | Gly | Thr | Phe | Thr | Ser | Asp | Xaa | Ser |
|    | 18  | 19  | 20  | 21  | 22  | 23  | 24  | 25  | 26  | 27  | 28  |
|    | Xaa | Xaa | Xaa | Glu | Xaa | Xaa | Ala | Xaa | Xaa | Xaa | Phe |
| 25 | 29  | 30  | 31  | 32  | 33  | 34  | 35  | 36  | 37  |     |     |
|    | Ile | Xaa | Trp | Leu | Xaa | Xaa | Gly | Xaa | R   |     |     |
- formula VIII (SEQ ID NO: 11)

wherein:

- 30 Xaa at position 7 is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine, homohistidine,  $\alpha$ -fluoromethyl-histidine or  $\alpha$ -methyl-histidine;
- Xaa at position 8 is: Gly, Ala, or Val;
- 35 Xaa at position 16 is: Leu or Val;

- 112 -

- Xaa at position 18 is Lys or Ser;  
Xaa at position 19 is: Gln or Tyr;  
Xaa at position 20 is: Met or Leu;  
Xaa at position 22 is: Glu or Gln;  
5 Xaa at position 23 is: Glu, or Gln;  
Xaa at position 25 is: Val or Ala;  
Xaa at position 26 is: Arg or Lys;  
Xaa at position 27 is Leu or Glu;  
Xaa at position 30 is: Glu or Ala;  
10 Xaa at position 33 is: Val or Lys;  
Xaa at position 34 is: Asn or Lys;  
Xaa at position 36 is: Gly or Arg; and  
Xaa at position 37 is: Gly, Pro, Pro-Ser-Ser-Gly-Ala-Pro-  
Pro-Pro-Ser, or is absent.
- 15 36. The heterologous fusion protein of claims 27 through 35 wherein the GLP-1 compound has no more than 6 amino acids that are different from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4.
- 20 37. The heterologous fusion protein of Claim 36 wherein the GLP-1 compound has no more than 5 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4.
- 25 38. The heterologous fusion protein of Claim 37 wherein the GLP-1 compound has no more than 4 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4.
- 30 39. The heterologous fusion protein of Claim 38 wherein the GLP-1 compound has no more than 3 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4.

35

- 113 -

40. The heterologous fusion protein of Claim 39 wherein the GLP-1 compound has no more than 2 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH, GLP-1(7-36)OH, or Exendin-4.

5

41. The heterologous fusion protein of any one of Claims 27 through 40 wherein Xaa at position 8 is glycine or valine.

42. The heterologous fusion protein of Claim 34 wherein the GLP-1 compound has no more than 2 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH or GLP-1(7-36)OH and Xaa at position 8 is glycine or valine and Xaa at position 30 is alanine, glutamic acid, aspartic acid, serine, or histidine.

15

43. The heterologous fusion protein of Claim 42 wherein Xaa at position 30 is Glutamic acid.

44. The heterologous fusion protein of Claim 34 wherein the GLP-1 compound has no more than 2 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH or GLP-1(7-36)OH and Xaa at position 8 is glycine or valine and Xaa at position 37 is histidine, phenylalanine, tyrosine, tryptophan, aspartic acid, glutamic acid, serine, threonine, arginine, or lysine.

20

45. The heterologous fusion protein of Claim 44 wherein Xaa at position 37 is histidine.

25

- 114 -

46. The heterologous fusion protein of Claim 34 wherein the GLP-1 compound has no more than 2 amino acids that differ from the corresponding amino acid in GLP-1(7-37)OH or GLP-1(7-36)OH and Xaa at position 8 is glycine, valine, leucine, isoleucine, serine, threonine, or methionine and Xaa at position 22 is aspartic acid, glutamic acid, lysine, arginine, asparagine, glutamine or histidine.
47. The heterologous fusion protein of Claim 46 wherein Xaa at position 22 is lysine or glutamic acid.
48. The heterologous fusion protein of Claim 34 wherein the GLP-1 compound is Val-8-GLP-1(7-37).
49. The heterologous fusion protein of Claim 34 wherein the GLP-1 compound is Gly-8-GLP-1(7-37).
50. The heterologous fusion protein of any one of Claims 27 through 49 wherein the second polypeptide is the Fc portion of an Ig selected from the group consisting of: IgG1, IgG2, IgG3, IgG4, IgE, IgA, IgD, or IgM.
51. The heterologous fusion protein of any one of Claims 27 through 50 wherein the second polypeptide is the Fc portion of an Ig selected from the group consisting of: IgG1, IgG2, IgG3, and IgG4.
52. The heterologous fusion protein of Claim 51 wherein the second polypeptide is the Fc portion of an IgG1 immunoglobulin.
53. The heterologous fusion protein of Claim 51 wherein the second polypeptide is the Fc portion of an IgG4 immunoglobulin.

- 115 -

54. The heterologous fusion protein of Claims 50 through 53 wherein the IgG is human.
- 5 55. The heterologous fusion protein of any one of Claims 27 through 54 wherein the Fc portion comprises the hinge, CH2, and CH3 domains.
56. The heterologous fusion protein of Claim 52 wherein the  
10 second polypeptide has the sequence of SEQ ID NO: 32.
57. A polynucleotide encoding a heterologous fusion protein of any one of Claims 1 through 56.
- 15 58. A vector comprising the polynucleotide of Claim 57.
59. A host cell comprising the vector of Claim 58.
60. A host cell expressing at least one heterologous fusion  
20 protein of any one of Claims 1 through 56.
61. The host cell of Claim 60 wherein said host cell is a CHO cell.
- 25 62. A process for producing a heterologous fusion protein comprising the steps of transcribing and translating a polynucleotide of Claim 57 under conditions wherein the heterologous fusion protein is expressed in detectable amounts.
- 30 63. A method for normalizing blood glucose levels in a mammal in need thereof comprising the administration of a therapeutically effective amount of the heterologous fusion protein of any one of Claims 1 through 56.

35



- 116 -

64. A method of treating a patient with non-insulin dependent diabetes mellitus comprising the administration of a therapeutically effective amount of the heterologous fusion protein of any one of Claims 1 through 56.

5

65. A method of treating obesity comprising the administration of a therapeutically effective amount of the heterologous fusion protein of any one of Claims 1 through 56.

10

66. The use of a heterologous fusion protein as claimed in any one of Claims 1 through 56 for the manufacture of a medicament for the treatment of patients with non-insulin dependent diabetes mellitus.

15

67. The use of a heterologous fusion protein as claimed in any one of Claims 1 through 56 for the manufacture of a medicament for the treatment of patients with obesity.

20

68. A pharmaceutical formulation adapted for the treatment of patients with non-insulin dependent diabetes comprising a heterologous fusion protein of any one of Claims 1 through 56.

25

69. The heterologous fusion protein of Claims 1, 2, or 3 wherein the GLP-1 compound comprises the sequence of formula IX [SEQ ID NO: 12]

30

Xaa<sub>7</sub>-Xaa<sub>8</sub>-Glu-Gly-Thr-Xaa<sub>12</sub>-Thr-Ser-Asp-Xaa<sub>16</sub>-Ser-  
Xaa<sub>18</sub>-Xaa<sub>19</sub>-Xaa<sub>20</sub>-Glu-Xaa<sub>22</sub>-Gln-Ala-Xaa<sub>25</sub>-Lys-Xaa<sub>27</sub>-  
Phe-Ile-Xaa<sub>30</sub>-Trp-Leu-Xaa<sub>33</sub>-Lys-Gly-Arg-Xaa<sub>37</sub>

Formula IX (SEQ ID NO: 12)

wherein:

35 Xaa<sub>7</sub> is: L-histidine, D-histidine, desamino-histidine, 2-

- 117 -

amino-histidine,  $\beta$ -hydroxy-histidine,  
homohistidine,  $\alpha$ -fluoromethyl-histidine, or  $\alpha$ -  
methyl-histidine;

- Xaa<sub>8</sub> is: Ala, Gly, Val, Leu, Ile, Ser, or Thr;  
5 Xaa<sub>12</sub> is: Phe, Trp, or Tyr;  
Xaa<sub>15</sub> is: Val, Trp, Ile, Leu, Phe, or Tyr;  
Xaa<sub>18</sub> is: Ser, Trp, Tyr, Phe, Lys, Ile, Leu, Val;  
Xaa<sub>19</sub> is: Tyr, Trp, or Phe;  
Xaa<sub>20</sub> is: Leu, Phe, Tyr, or Trp;  
10 Xaa<sub>22</sub> is: Gly, Glu, Asp, or Lys;  
Xaa<sub>25</sub> is: Ala, Val, Ile, or Leu;  
Xaa<sub>27</sub> is: Glu, Ile, or Ala;  
Xaa<sub>30</sub> is: Ala or Glu  
Xaa<sub>33</sub> is: Val, or Ile; and  
15 Xaa<sub>37</sub> is: Gly, His, NH<sub>2</sub>, or is absent.

70. The heterologous fusion protein of Claim 69 wherein the  
second polypeptide is human albumin.

- 20 71. The heterologous fusion protein of Claim 70 wherein the  
second polypeptide has the sequence of SEQ ID NO: 34.

72. The heterologous fusion protein of Claim 69 wherein the  
second polypeptide is an N-terminal fragment of albumin.

25

73. The heterologous fusion protein of any one of Claims  
27, 28, or 29 wherein the GLP-1 compound comprises the  
sequence of formula IX [SEQ ID NO: 12]

- 30 Xaa<sub>7</sub>-Xaa<sub>8</sub>-Glu-Gly-Thr-Xaa<sub>12</sub>-Thr-Ser-Asp-Xaa<sub>16</sub>-Ser-  
Xaa<sub>18</sub>-Xaa<sub>19</sub>-Xaa<sub>29</sub>-Glu-Xaa<sub>22</sub>-Gln-Ala-Xaa<sub>25</sub>-Lys-Xaa<sub>27</sub>-  
Phe-Ile-Xaa<sub>30</sub>-Trp-Leu-Xaa<sub>33</sub>-Lys-Gly-Arg-Xaa<sub>37</sub>

Formula IX (SEQ ID NO: 12)

- 118 -

wherein:

- Xaa<sub>7</sub> is: L-histidine, D-histidine, desamino-histidine, 2-amino-histidine,  $\beta$ -hydroxy-histidine,  
5 homohistidine,  $\alpha$ -fluoromethyl-histidine, or  $\alpha$ -methyl-histidine;
- Xaa<sub>8</sub> is: Ala, Gly, Val, Leu, Ile, Ser, or Thr;  
Xaa<sub>12</sub> is: Phe, Trp, or Tyr;  
Xaa<sub>16</sub> is: Val, Trp, Ile, Leu, Phe, or Tyr;  
10 Xaa<sub>18</sub> is: Ser, Trp, Tyr, Phe, Lys, Ile, Leu, Val;  
Xaa<sub>19</sub> is: Tyr, Trp, or Phe;  
Xaa<sub>20</sub> is: Leu, Phe, Tyr, or Trp;  
Xaa<sub>22</sub> is: Gly, Glu, Asp, or Lys;  
Xaa<sub>25</sub> is: Ala, Val, Ile, or Leu;  
15 Xaa<sub>27</sub> is: Glu, Ile, or Ala;  
Xaa<sub>30</sub> is: Ala or Glu  
Xaa<sub>33</sub> is: Val, or Ile; and  
Xaa<sub>37</sub> is: Gly, His, NH<sub>2</sub>, or is absent.
- 20 74. The heterologous fusion protein of Claim 73 wherein the second polypeptide is the Fc portion of an Ig selected from the group consisting of: IgG1, IgG2, IgG3, IgG4, IgE, IgA, IgD, or IgM.
- 25 75. The heterologous fusion protein of Claim 73 or 74 wherein the second polypeptide is the Fc portion of an Ig selected from the group consisting of: IgG1, IgG2, IgG3, and IgG4.
- 30 76. The heterologous fusion protein of Claim 75 wherein the second polypeptide is the Fc portion of an IgG1 immunoglobulin.

- 119 -

77. The heterologous fusion protein of Claim 75 wherein the second polypeptide is the Fc portion of an IgG4 immunoglobulin.
- 5 78. The heterologous fusion protein of Claims 73 through 78 wherein the IgG is human.
79. The heterologous fusion protein of Claim 75 wherein the second polypeptide has the sequence of SEQ ID NO: 32.
- 10 80. The heterologous fusion protein of any one of Claims 69 through 79 wherein the GLP-1 compound is selected from the group consisting of Gly<sup>8</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>12</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>12</sup>-GLP-1(7-36), Val<sup>8</sup>-Trp<sup>12</sup>-GLP-1(7-37), Val<sup>8</sup>-Leu<sup>16</sup>-GLP-1(7-37), Val<sup>8</sup>-Val<sup>16</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>16</sup>-GLP-1(7-37), Gly<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Leu<sup>25</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>12</sup>-Tyr<sup>16</sup>-GLP-1(7-37), Val<sup>8</sup>-Trp<sup>12</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>12</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>16</sup>-Phe<sup>19</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Trp<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Leu<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Ile<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Phe<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Trp<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Phe<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Ile<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Lys<sup>18</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Trp<sup>19</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Phe<sup>19</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Phe<sup>20</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-Leu<sup>25</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-Ile<sup>25</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-Val<sup>25</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-Ile<sup>27</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-Ala<sup>27</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-Ile<sup>33</sup>-GLP-1(7-37), Val<sup>8</sup>-Asp<sup>6</sup>-Ile<sup>11</sup>-Tyr<sup>16</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Tyr<sup>16</sup>-Trp<sup>19</sup>-Glu<sup>22</sup>-GLP-1(7-37), Val<sup>8</sup>-Trp<sup>16</sup>-Glu<sup>22</sup>-Val<sup>25</sup>-Ile<sup>33</sup>-GLP-1(7-37), Val<sup>8</sup>-Trp<sup>16</sup>-Glu<sup>22</sup>-Ile<sup>33</sup>-GLP-1(7-37), Val<sup>8</sup>-Glu<sup>22</sup>-Val<sup>25</sup>-Ile<sup>33</sup>-GLP-1(7-37), Val<sup>8</sup>-Trp<sup>16</sup>-

- 120 -

Glu<sup>22</sup>-Val<sup>25</sup>- GLP-1(7-37), and Val<sup>8</sup>-Cys<sup>16</sup>-Lys<sup>26</sup>-GLP-1(7-37).

1/17

Fig. 1

				5					10					15
Ala	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys
				20						25				30
Pro	Ala	Pro	Glu	Lys	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro
				35						40				45
Lys	Pro	Lys	Asp	Thr	Lys	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
				50						55				60
Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe
				65						70				75
Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys
				80						85				90
Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val
				95						100				105
Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
				110						115				120
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr
				125						130				135
Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr
				140						145				150
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu
				155						160				165
Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu
				170						175				180
Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro
				185						190				195
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu
				200						205				210
Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys
				215						220				225
Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser
				230										
Leu	Ser	Leu	Ser	Pro	Gly	Lys								

[SEQ ID NO: 32]

2/17

Fig. 2


					5					10					15				
Asp	Ala	His	Lys	Ser	Glu	Val	Ala	His	Arg	Phe	Lys	Asp	Leu	Gly	Glu	Glu	Asn		
20					25						30					35			
Phe	Lys	Ala	Leu	Val	Leu	Ile	Ala	Phe	Ala	Gln	Tyr	Leu	Gln	Gln	Cys	Pro	Phe		
			40					45					50						
Glu	Asp	His	Val	Lys	Leu	Val	Asn	Glu	Val	Thr	Glu	Phe	Ala	Lys	Thr	Cys	Val		
55					60					65					70				
Ala	Asp	Glu	Ser	Ala	Glu	Asn	Cys	Asp	Lys	Ser	Leu	His	Thr	Leu	Phe	Gly	Asp		
			75				80					85				90			
Lys	Leu	Cys	Thr	Val	Ala	Thr	Leu	Arg	Glu	Thr	Tyr	Gly	Glu	Met	Ala	Asp	Cys		
				95					100					105					
Cys	Ala	Lys	Gln	Glu	Pro	Glu	Arg	Asn	Glu	Cys	Phe	Leu	Gln	His	Lys	Asp	Asp		
	110					115					120					125			
Asn	Pro	Asn	Leu	Pro	Arg	Leu	Val	Arg	Pro	Glu	Val	Asp	Val	Met	Cys	Thr	Ala		
			130					135					140						
Phe	His	Asp	Asn	Glu	Glu	Thr	Phe	Leu	Lys	Lys	Tyr	Leu	Tyr	Glu	Ile	Ala	Arg		
145					150					155					160				
Arg	His	Pro	Tyr	Phe	Tyr	Ala	Pro	Glu	Leu	Leu	Phe	Phe	Ala	Lys	Arg	Tyr	Lys		
		165				170					175					180			
Ala	Ala	Phe	Thr	Glu	Cys	Cys	Gln	Ala	Ala	Asp	Lys	Ala	Ala	Cys	Leu	Leu	Pro		
				185					190					195					
Lys	Leu	Asp	Glu	Leu	Arg	Asp	Glu	Gly	Lys	Ala	Ser	Ser	Ala	Lys	Gln	Arg	Leu		
	200					205					210					215			
Lys	Cys	Ala	Ser	Leu	Gln	Lys	Phe	Gly	Glu	Arg	Ala	Phe	Lys	Ala	Trp	Ala	Val		
			220				225						230						
Ala	Arg	Leu	Ser	Gln	Arg	Phe	Pro	Lys	Ala	Glu	Phe	Ala	Glu	Val	Ser	Lys	Leu		
235					240					245					250				
Val	Thr	Asp	Leu	Thr	Lys	Val	His	Thr	Glu	Cys	Cys	His	Gly	Asp	Leu	Leu	Glu		
		255					260					265				270			
Cys	Ala	Asp	Asp	Arg	Ala	Asp	Leu	Ala	Lys	Tyr	Ile	Cys	Glu	Asn	Gln	Asp	Ser		
				275					280					285					
Ile	Ser	Ser	Lys	Leu	Lys	Glu	Cys	Cys	Glu	Lys	Pro	Leu	Leu	Glu	Lys	Ser	His		
	290					295					300					305			
Cys	Ile	Ala	Glu	Val	Glu	Asn	Asp	Glu	Met	Pro	Ala	Asp	Leu	Pro	Ser	Leu	Ala		
			310					315					320						
Ala	Asp	Phe	Val	Glu	Ser	Lys	Asp	Val	Cys	Lys	Asn	Tyr	Ala	Glu	Ala	Lys	Asp		
325					330					335					340				
Val	Phe	Leu	Gly	Met	Phe	Leu	Tyr	Glu	Tyr	Ala	Arg	Arg	His	Pro	Asp	Tyr	Ser		
		345					350				355					360			
Val	Val	Leu	Leu	Leu	Arg	Leu	Ala	Lys	Thr	Tyr	Glu	Thr	Thr	Leu	Glu	Lys	Cys		
				365					370					375					
Cys	Ala	Ala	Ala	Asp	Pro	His	Glu	Cys	Tyr	Ala	Lys	Val	Phe	Asp	Glu	Phe	Lys		
	380					385					390					395			
Pro	Leu	Val	Glu	Glu	Pro	Gln	Asn	Leu	Ile	Lys	Gln	Asn	Cys	Glu	Leu	Phe	Glu		
			400					405					410						
Asn	Leu	Gly	Glu	Tyr	Lys	Phe	Gln	Asn	Ala	Leu	Leu	Val	Arg	Tyr	Thr	Lys	Lys		
415					420					425					430				
Val	Pro	Gln	Val	Ser	Thr	Pro	Thr	Leu	Val	Glu	Val	Ser	Arg	Asn	Leu	Gly	Lys		
		435					440					445				450			
Val	Gly	Ser	Lys	Cys	Cys	Lys	His	Pro	Glu	Ala	Lys	Arg	Met	Pro	Cys	Ala	Glu		
				455					460					465					
Asp	Tyr	Leu	Ser	Val	Val	Leu	Asn	Gln	Leu	Cys	Val	Leu	His	Glu	Lys	Thr	Pro		
	470					475					480					485			
Val	Ser	Asp	Arg	Val	Thr	Lys	Cys	Cys	Thr	Glu	Ser	Leu	Val	Asn	Arg	Arg	Pro		






4/17

Fig. 3




5/17

Fig. 4




6/17

Fig. 5




7/17

Fig. 6




8/17

Fig. 7




9/17

Fig. 8



10/17


Fig. 9



11/17

Fig. 10

A.






12/17

Fig. 10


B.



13/17

Fig. 11


A.



14/17

Fig. 11

B.



15/17

Fig. 12

1  
GAGCCCCAATCTTGTGACAAACTCACACATGCCACCCTGCCAGCACC  
50  
TGAACCTCTGGGGGACCGTCAGTCTTCCTCTTCCCCCAAACCCAAGG  
100  
ACACCTCATGATCTCCCGACCCCTGAGGTACATGCGTGGTGGTGGAC  
150  
GTGAGCCACGAAGACCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGT  
200  
GGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCA  
250  
CGTACCGTGTGGTCAGCGTCTCACCGTCTTGCAACAGGACTGGCTGAAT  
300  
GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCAT  
350  
CGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCAAGGTGT  
400  
ACACCTTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGAGCCTG  
450  
ACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGA  
500  
GAGCAATGGGCAGCCGGAGAACAACACTACAGACCACGCCTCCCGTGTGG  
550  
ACTCCGACGGCTCCTTCTTCTCTATAGCAAGCTCACCGTGGACAAGAGC  
600  
AGGTGGCAGCAGGGGAACGTCTTCTCATGTCCGTGATGATGAGGCTCT  
650  
GCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGAT  
700  
AGT [SEQ ID NO: 33]

16/17

Fig. 13

GATGCGCACAAGAGTGAGGTTGCTCATCGGTTTAAAGATTGGGAGAAGAA  
 50  
 AAATTTCAAAGCCCTGGTGTGATTGCCTTTGCTCAGTATCTTCAGCAGT  
 100  
 GTCCATTGGAAGATCATGTAAATTAGTGAATGAAGTAACTGAATTTGCA  
 150  
 AAAACATGTGTTGCTGATGAGTCAGCTGAAAATTTGTGACAAATCACTTCA  
 200  
 TACCCTTTTGGAGACAAATTATGCACAGTTGCAACTCTTCGTGAAACCT  
 250  
 ATGGTGAAATGGCTGACTGCTGTGCAAAACAAGAACCTGAGAGAAATGAA  
 300  
 TGCTTCTTGCAACACAAGATGACAACCCAAACCTCCCCCGATTGGTGAG  
 350  
 ACCCAGAGGTTGATGTGATGTGCACTGCTTTTCATGACAAATGAAGAGACAT  
 400  
 TTTTGAAAAATACTTATATGAAATTGCCAGAAGACATCCTTACTTTTAT  
 450  
 GCCCCGGAACCTCCTTTCTTTTGCTAAAGGTTATAAAGCTGCTTTTACAGA  
 500  
 ATGTTGCCAAGCTGCTGATAAAGCTGCCTGCCTGTTGCCAAAGCTCGATG  
 550  
 AACCTTCGGGATGAAGGGAAGGCTTCCTCTGCCAAACAGAGACTCAAGTGT  
 600  
 GCCAGCTCTCCAAAAATTTGGAGAAAAGAGCTTTCAAAGCATGGGCGATAGC  
 650  
 TCGCCTGAGCCAGAGATTCCCAAAGCTGAGTTTGCAGAAGTTTCCAAGT  
 700  
 TAGTGACAGATCTTTACCAAAGTCCACACGGAATGCTGCCATGGAGATCTG  
 750  
 CTTGGAATGTGCTGATGACAGGGCGGACCTTGCCAAGTATATCTGTGAAAA  
 800  
 TCAAGATTTCGATCTCCAGTAACTGAAGGAATGCTGTGAAAAAACCTCTGT  
 850  
 TGGAAAAATCCCACTGCACTGTGCCGAAGTGGAAAAATGATGAGATGCCTGCT  
 900  
 GACTTGCCTTCATTAGCTGCTGATTTTGTGTGAAAGTAAGGATGTTTGCAA  
 950  
 AAACATATGCTGAGGCAAAGGATGTCTTCTCTGGGCATGTTTTTGTATGAAT  
 1000  
 ATGCAAGAAGGCATCCTGATTACTCTGTCTGTCTGCTGCTGCTGAGACTTGCC

17/17

## Fig. 13 Continued

1050  
AAGACATATGAAACCACTCTAGAGAAGTGCTGTGCCGCTGCAGATCCTCA  
1100  
TGAATGCTATGCCAAAGTGTTGATGAATTTAAACCTCTTGTGGAAGAGC  
1150  
CTCAGAATTTAATCAAACAAAATTGTGAGCTTTTTGAGCAGCTTGGAGAG  
1200  
TACAAATTCAGAATGCGCTATTAGTTCGTTACACCAAGAAAGTACCCCA  
1250  
AGTGTCAACTCCAACCTTTGTAGAGGCTCAAGAAACCTAGGAAAAGTGG  
1300  
GCAGCAAATGTTGTAAACATCCTGAAGCAAAAAGAATGCCCTGTGCAGAA  
1350  
GACTATCTATCCGTGGTCCTGAACCAGTTATGTGTGTTGCATGAGAAAAC  
1400  
GCCAGTAAAGTGACAGAGTCACCAAATGCTGCACAGAATCCCTTGGTGAACA  
1450  
GGCGACCATGCTTTTCAGCTCTGGAAGTCGATGAAACATACGTTCCCAA  
1500  
GAGTTTAATGCTGAAACATTACCTTCCATGCAGATATATGCACACTTTC  
1550  
TGAGAAAGGAGAGACAAATCAAGAAACAAACTGCACCTTGTTGAGCTCGTGA  
1600  
AACACAAGCCCAAGGCAACAAAAGAGCAACTGAAAGCTGTTATGGATGAT  
1650  
TTGCGAGCTTTTGTAGAGAAGTGCTGCAAGGCTGACGATAAGGAGACCTG  
1700  
CTTTGCCGAGGAGGGTAAAAAACTTGTGTGTCAGATCAAGCTGCCTTAG  
1750  
GCTTATAATGAC           [SEQ ID NO: 35]

X-13991.ST25.txt  
SEQUENCE LISTING

&lt;110&gt; Eli Lilly and Company

&lt;120&gt; GLP-1 FUSION PROTEINS

&lt;130&gt; X-13991

&lt;150&gt; US 60/251,954

&lt;151&gt; 2000-06-12

&lt;160&gt; 35

&lt;170&gt; PatentIn version 3.1

&lt;210&gt; 1

&lt;211&gt; 31

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 1

His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly  
1 5 10 15Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly  
20 25 30

&lt;210&gt; 2

&lt;211&gt; 39

&lt;212&gt; PRT

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

X-13991.ST25.txt

&lt;222&gt; (2)..(2)

&lt;223&gt; Xaa at position 2 is Ala, Gly, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (3)..(3)

&lt;223&gt; Xaa at position 3 is Glu, Asp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (5)..(5)

&lt;223&gt; Xaa at position 5 is Thr, Ala, Gly, Ser, Leu, Ile, Val, Glu, Asp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (8)..(8)

&lt;223&gt; Xaa at position 8 is Ser, Ala, Gly, Thr, Leu, Ile, Val, Glu, Asp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (10)..(10)

&lt;223&gt; Xaa at position 10 is Val, Ala, Gly, Ser, Thr, Leu, Ile, Tyr, Glu, Asp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (11)..(11)

&lt;223&gt; Xaa at position 11 is Ser, Ala, Gly, Thr, Leu, Ile, Val, Glu, Asp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (12)..(12)



X-13991.ST25.txt

<223> Xaa at position 12 is Ser, Ala, Gly, Thr, Leu, Ile, Val, Glu, Asp, Lys, Trp or Tyr;

<220>

<221> MISC\_FEATURE

<222> (13)..(13)

<223> Xaa at position 13 is Tyr, Phe, Trp, Glu, Asp, Gln, or Lys;

<220>

<221> MISC\_FEATURE

<222> (14)..(14)

<223> Xaa at position 14 is Leu, Ala, Gly, Ser, Thr, Ile, Val, Glu, Asp, Met, Lys, Trp or Tyr;

<220>

<221> MISC\_FEATURE

<222> (15)..(15)

<223> Xaa at position 15 is Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (16)..(16)

<223> Xaa at position 16 is Gly, Ala, Ser, Thr, Leu, Ile, Val, Glu, Asp, Trp or Lys;

<220>

<221> MISC\_FEATURE

<222> (17)..(17)

<223> Xaa at position 17 is Gln, Asn, Arg, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (18)..(18)

X-13991.ST25.txt

<223> Xaa at position 18 is Ala, Gly, Ser, Thr, Leu, Ile, Val, Arg, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (19)..(19)

<223> Xaa at position 19 is Ala, Gly, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (20)..(20)

<223> Xaa at position 20 is Lys, Arg, Gln, Glu, Asp, or His;

<220>

<221> MISC\_FEATURE

<222> (21)..(21)

<223> Xaa at position 21 is Leu, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (24)..(24)

<223> Xaa at position 24 is Ala, Gly, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (25)..(25)

<223> Xaa at position 25 is Trp, Phe, Tyr, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (26)..(26)

X-13991.ST25.txt

<223> Xaa at position 26 is Leu, Gly, Ala, Ser, Thr, Ile, Val, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (27)..(27)

<223> Xaa at position 27 is Val, Gly, Ala, Ser, Thr, Leu, Ile, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (28)..(28)

<223> Xaa at position 28 is Asn, Lys, Arg, Glu, Asp, or His;

<220>

<221> MISC\_FEATURE

<222> (29)..(29)

<223> Xaa at position 29 is Gly, Ala, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys;

<220>

<221> MISC\_FEATURE

<222> (30)..(30)

<223> Xaa at position 30 is Gly, Arg, Lys, Glu, Asp, or His;

<220>

<221> MISC\_FEATURE

<222> (31)..(31)

<223> Xaa at position 31 is Pro, Gly, Ala, Ser, Thr, Leu, Ile, Val, Glu, Asp, or Lys, or is deleted;

<220>

<221> MISC\_FEATURE

X-13991.ST25.txt

&lt;222&gt; (32)..(32)

&lt;223&gt; Xaa at position 32 is Ser, Arg, Lys, Glu, Asp, or His, or is deleted;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (33)..(33)

&lt;223&gt; Xaa at position 33 is Ser, Arg, Lys, Glu, Asp, or His, or is deleted;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (34)..(34)

&lt;223&gt; Xaa at position 34 is Gly, Asp, Glu, or Lys, or is deleted;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (35)..(35)

&lt;223&gt; Xaa at position 35 is Ala, Phe, Trp, Tyr, Glu, Asp, or Lys, or is deleted;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (36)..(36)

&lt;223&gt; Xaa at position 36 is Ser, Pro, Lys, Glu, or Asp, or is deleted;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (37)..(37)

&lt;223&gt; Xaa at position 37 is Ser, Pro, Glu, Asp, or Lys, or is deleted;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

X-13991.ST25.txt

&lt;222&gt; (38)..(38)

&lt;223&gt; Xaa at position 38 is Gly, Pro, Glu, Asp, or Lys, or is deleted;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (39)..(39)

&lt;223&gt; Xaa at position 39 is Ala, Ser, Val, Glu, Asp, or Lys, or is deleted;

&lt;400&gt; 2

His	Xaa	Xaa	Gly	Xaa	Phe	Thr	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1			5					10					15		

Xaa	Xaa	Xaa	Xaa	Xaa	Phe	Ile	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
			20				25						30		

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
			35				

&lt;210&gt; 3

&lt;211&gt; 32

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (1)..(1)

&lt;223&gt; Xaa at position 1 is L-histidine, D-histidine, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (2)..(2)

&lt;223&gt; Xaa at position 2 is Gly, Ala, Val, Leu, Ile, Ser, or Thr;

&lt;220&gt;

X-13991.ST25.txt

<221> MISC\_FEATURE  
<222> (3)..(3)  
<223> Xaa at position 3 is Thr, Ser, Arg, Lys, Trp, Phe, Tyr, Glu, or His;

<220>  
<221> MISC\_FEATURE  
<222> (5)..(5)  
<223> Xaa at position 5 is Asp, Glu, Arg, Thr, Ala, Lys, or His;

<220>  
<221> MISC\_FEATURE  
<222> (6)..(6)  
<223> Xaa at position 6 is His, Trp, Phe, or Tyr;

<220>  
<221> MISC\_FEATURE  
<222> (10)..(10)  
<223> Xaa at position 10 is Leu, Ser, Thr, Trp, His, Phe, Asp, Val, Tyr, Glu, or Ala;

<220>  
<221> MISC\_FEATURE  
<222> (12)..(12)  
<223> Xaa at position 12 is His, Pro, Asp, Glu, Arg, Ser, Ala, or Lys;

<220>  
<221> MISC\_FEATURE  
<222> (13)..(13)  
<223> Xaa at position 13 is Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;

<220>

X-13991.ST25.txt

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (17)..(17)

&lt;223&gt; Xaa at position 17 is His, Asp, Lys, Glu, Gln, or Arg;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (18)..(18)

&lt;223&gt; Xaa at position 18 is Glu, Arg, Ala, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (20)..(20)

&lt;223&gt; Xaa at position 20 is Trp, Tyr, Phe, Asp, Lys, Glu, or His;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (21)..(21)

&lt;223&gt; Xaa at position 21 is Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (24)..(24)

&lt;223&gt; Xaa at position 24 is Ala, Glu, Asp, Ser, or His;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (25)..(25)

&lt;223&gt; Xaa at position 25 is Asp, Glu, Ser, Thr, Arg, Trp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

X-13991.ST25.txt

&lt;222&gt; (27)..(27)

&lt;223&gt; Xaa at position 27 is Asp, Arg, Val, Lys, Ala, Gly, or Glu;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (28)..(28)

&lt;223&gt; Xaa at position 28 is Glu, Lys, or Asp;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (29)..(29)

&lt;223&gt; Xaa at position 29 is Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (30)..(30)

&lt;223&gt; Xaa at position 30 is Thr, Ser, Asp, Trp, Tyr, Phe, Arg, Glu, or His;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (31)..(31)

&lt;223&gt; Xaa at position 31 is Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (32)..(32)

&lt;223&gt; Xaa at position 31 is Pro or is deleted.

&lt;400&gt; 3

Xaa	Xaa	Xaa	Gly	Xaa	Xaa	Thr	Ser	Asp	Xaa	Ser	Xaa	Xaa	Leu	Glu	Gly
1			5						10					15	

Xaa	Xaa	Ala	Xaa	Xaa	Phe	Ile	Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
		20					25						30		



X-13991.ST25.txt

<210> 4  
<211> 32  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> synthetic construct  
<220>  
<221> MISC\_FEATURE  
<222> (1)..(1)  
<223> Xaa at position 1 is L-histidine, D-histidine, or is deleted.

<220>  
<221> MISC\_FEATURE  
<222> (2)..(2)  
<223> Xaa at position 2 is Gly, Ala, Val, Leu, Ile, Ser, or Thr;

<220>  
<221> MISC\_FEATURE  
<222> (5)..(5)  
<223> Xaa at position 5 is Asp, Glu, Arg, Thr, Ala, Lys, or His;

<220>  
<221> MISC\_FEATURE  
<222> (6)..(6)  
<223> Xaa at position 6 is His, Trp, Phe, or Tyr;

<220>  
<221> MISC\_FEATURE  
<222> (10)..(10)  
<223> Xaa at position 10 is Leu, Ser, Thr, Trp, His, Phe, Asp, Val, Glu, or Ala;

X-13991.ST25.txt

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (16)..(16)

&lt;223&gt; Xaa at position 16 is Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (17)..(17)

&lt;223&gt; Xaa at position 17 is His, Asp, Lys, Glu, or Gln;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (18)..(18)

&lt;223&gt; Xaa at position 18 is Glu, His, Ala, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (19)..(19)

&lt;223&gt; Xaa at position 19 is Asp, Lys, Glu, or His;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (21)..(21)

&lt;223&gt; Xaa at position 21 is Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (24)..(24)

&lt;223&gt; Xaa at position 24 is Ala, Glu, Asp, Ser, or His;

X-13991.ST25.txt

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (27)..(27)

&lt;223&gt; Xaa at position 27 is Asp, Arg, Val, Lys, Ala, Gly, or Glu;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (28)..(28)

&lt;223&gt; Xaa at position 28 is Glu, Lys, or Asp;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (29)..(29)

&lt;223&gt; Xaa at position 29 is Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (30)..(30)

&lt;223&gt; Xaa at position 30 is Arg, Glu, or His;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (31)..(31)

&lt;223&gt; Xaa at position 31 is Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (32)..(32)

&lt;223&gt; Xaa at position 32 is Pro, or is deleted.

&lt;400&gt; 4

Xaa Xaa Glu Gly Xaa Xaa Thr Ser Asp Xaa Ser Ser Tyr Leu Glu Xaa  
 1 5 10 15

x-13991.ST25.txt

Xaa Xaa Xaa Lys Xaa Phe Ile Xaa Trp Leu Xaa Xaa Xaa Xaa Xaa Xaa  
20 25 30

&lt;210&gt; 5

&lt;211&gt; 32

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (1)..(1)

&lt;223&gt; Xaa at position 1 is L-histidine, D-histidine, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (2)..(2)

&lt;223&gt; Xaa at position 2 is Gly, Ala, Val, Leu, Ile, Ser, Met, or Thr;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (6)..(6)

&lt;223&gt; Xaa at position 6 is His, Trp, Phe, or Tyr;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (10)..(10)

&lt;223&gt; Xaa at position 10 is Leu, Ser, Thr, Trp, His, Phe, Asp, Val, Glu, or Ala;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (16)..(16)

X-13991.ST25.txt

<223> Xaa at position 16 is Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;

<220>

<221> MISC\_FEATURE

<222> (17)..(17)

<223> Xaa at position 17 is His, Asp, Lys, Glu, or Gln;

<220>

<221> MISC\_FEATURE

<222> (20)..(20)

<223> Xaa at position 20 is Asp, Lys, Glu, or His;

<220>

<221> MISC\_FEATURE

<222> (24)..(24)

<223> Xaa at position 24 is Ala, Glu, Asp, Ser, or His;

<220>

<221> MISC\_FEATURE

<222> (29)..(29)

<223> Xaa at position 29 is Thr, Ser, Lys, Arg, Trp, Tyr, Phe, Asp, Gly, Pro, His, or Glu;

<220>

<221> MISC\_FEATURE

<222> (31)..(31)

<223> Xaa at position 31 is Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, or is deleted.

<220>

<221> MISC\_FEATURE

<222> (32)..(32)

<223> Xaa at position 32 is Pro or is deleted.

X-13991.ST25.txt

&lt;400&gt; 5

Xaa Xaa Glu Gly Thr Xaa Thr Ser Asp Xaa Ser Ser Tyr Leu Glu Xaa  
 1 5 10 15

Xaa Ala Ala Xaa Glu Phe Ile Xaa Trp Leu Val Lys Xaa Arg Xaa Xaa  
 20 25 30

&lt;210&gt; 6

&lt;211&gt; 32

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (1)..(1)

&lt;223&gt; Xaa at position 1 is L-histidine, D-histidine, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (2)..(2)

&lt;223&gt; Xaa at position 2 is Gly, Ala, Val, Leu, Ile, Ser, or Thr;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (16)..(16)

&lt;223&gt; Xaa at position 16 is Gly, Asp, Glu, Gln, Asn, Lys, Arg, or Cys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (17)..(17)

&lt;223&gt; Xaa at position 17 is His, Asp, Lys, Glu, or Gln;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

X-13991.ST25.txt

&lt;222&gt; (18)..(18)

&lt;223&gt; Xaa at position 18 is Ala, Glu, His, Phe, Tyr, Trp, Arg, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (24)..(24)

&lt;223&gt; Xaa at position 24 is Ala, Glu, Asp, Ser, or His;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (31)..(31)

&lt;223&gt; Xaa at position 31 is Lys, Arg, Thr, Ser, Glu, Asp, Trp, Tyr, Phe, His, Gly, Gly-Pro, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (32)..(32)

&lt;223&gt; Xaa at position 32 is Pro or is deleted.

&lt;400&gt; 6

Xaa Xaa Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Xaa  
1 5 10 15Xaa Xaa Ala Lys Glu Phe Ile Xaa Trp Leu Val Lys Gly Arg Xaa Xaa  
20 25 30

&lt;210&gt; 7

&lt;211&gt; 31

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

X-13991.ST25.txt

&lt;222&gt; (1)..(1)

&lt;223&gt; Xaa at position 1 is L-histidine, D-histidine, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (2)..(2)

&lt;223&gt; Xaa at position 2 is Ala, Gly, Val, Thr, Ile, and alpha-methyl-Ala;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (15)..(15)

&lt;223&gt; Xaa at position 15 is Glu, Gln, Ala, Thr, Ser, and Gly;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (21)..(21)

&lt;223&gt; Xaa at position 21 is Glu, Gln, Ala, Thr, Ser, and Gly;

&lt;400&gt; 7

Xaa	Xaa	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Val	Ser	Ser	Tyr	Leu	Xaa	Gly
1			5					10						15	

Gln	Ala	Ala	Lys	Xaa	Phe	Ile	Ala	Trp	Leu	Val	Lys	Gly	Arg	Gly
		20					25						30	

&lt;210&gt; 8

&lt;211&gt; 30

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (19)..(19)



X-13991.ST25.txt  
 <223> Xaa at position 19 is Lys or Arg;

<220>

<221> MOD\_RES

<222> (27)..(27)

<223> ACETYLTATION

<220>

<221> MISC\_FEATURE

<222> (30)..(30)

<223> Xaa at position 30 is Gly;

<220>

<221> MOD\_RES

<222> (30)..(30)

<223> AMIDATION

<400> 8

Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln  
 1 5 10 15

Ala Ala Xaa Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Xaa  
 20 25 30

<210> 9

<211> 39

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 9

His Ser Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
 1 5 10 15

Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser  
 20 25 30

X-13991.ST25.txt

Ser Gly Ala Pro Pro Pro Ser  
35

&lt;210&gt; 10

&lt;211&gt; 39

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
1 5 10 15Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser  
20 25 30Ser Gly Ala Pro Pro Pro Ser  
35

&lt;210&gt; 11

&lt;211&gt; 39

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (1)..(1)

&lt;223&gt; Xaa at position 1 is L-histidine, D-histidine, or is deleted.

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (2)..(2)

&lt;223&gt; Xaa at position 2 is Gly, Ala, or Val;

&lt;220&gt;

X-13991.ST25.txt

<221> MISC\_FEATURE  
<222> (10)..(10)  
<223> Xaa at position 10 is Leu or Val;  
  
<220>  
<221> MISC\_FEATURE  
<222> (12)..(12)  
<223> Xaa at position 12 is Lys or Ser;  
  
<220>  
<221> MISC\_FEATURE  
<222> (13)..(13)  
<223> Xaa at position 13 is Gln or Tyr;  
  
<220>  
<221> MISC\_FEATURE  
<222> (14)..(14)  
<223> Xaa at position 14 is Met or Leu;  
  
<220>  
<221> MISC\_FEATURE  
<222> (16)..(16)  
<223> Xaa at position 16 is Glu or Gln;  
  
<220>  
<221> MISC\_FEATURE  
<222> (17)..(17)  
<223> Xaa at position 17 is Glu or Gln;  
  
<220>  
<221> MISC\_FEATURE  
<222> (19)..(19)  
<223> Xaa at position 19 is Val or Ala;

X-13991.ST25.txt

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (20)..(20)

&lt;223&gt; Xaa at position 20 is Arg or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (21)..(21)

&lt;223&gt; Xaa at position 21 is Leu or Glu;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (24)..(24)

&lt;223&gt; Xaa at position 24 is Glu or Ala;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (27)..(27)

&lt;223&gt; Xaa at position 27 is Val or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (28)..(28)

&lt;223&gt; Xaa at position 28 is Asn or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (30)..(30)

&lt;223&gt; Xaa at position 30 is Gly or Arg; and

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (31)..(31)

&lt;223&gt; Xaa at position 31 is Gly, or Pro;

X-13991.ST25.txt

<220>  
<221> MISC\_FEATURE  
<222> (32)..(32)  
<223> Xaa at position 32 is Ser, or is absent.

<220>  
<221> MISC\_FEATURE  
<222> (33)..(33)  
<223> Xaa at position 33 is Ser, or is absent.

<220>  
<221> MISC\_FEATURE  
<222> (34)..(34)  
<223> Xaa at position 34 is Gly, or is absent.

<220>  
<221> MISC\_FEATURE  
<222> (35)..(35)  
<223> Xaa at position 35 is Ala, or is absent.

<220>  
<221> MISC\_FEATURE  
<222> (36)..(36)  
<223> Xaa at position 36 is Pro, or is absent.

<220>  
<221> MISC\_FEATURE  
<222> (37)..(37)  
<223> Xaa at position 37 is Pro, or is absent.

<220>  
<221> MISC\_FEATURE  
<222> (38)..(38)

X-13991.ST25.txt  
 <223> Xaa at position 38 is Pro, or is absent.

<220>

<221> MISC\_FEATURE

<222> (39)..(39)

<223> Xaa at position 39 is Pro, or is absent.

<220>

<221> MISC\_FEATURE

<222> (39)..(39)

<223> Xaa at position 39 is Ser, or is absent.

<400> 11

Xaa Xaa Glu Glu Thr Phe Thr Ser Asp Xaa Ser Xaa Xaa Xaa Glu Xaa  
 1 5 10 15

Xaa Ala Xaa Xaa Xaa Phe Ile Xaa Trp Leu Xaa Xaa Gly Xaa Xaa Xaa  
 20 25 30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 35

<210> 12

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<220>

<221> MISC\_FEATURE

<222> (1)..(1)

<223> Xaa at position 1 is L-histidine, D-histidine, or is deleted.

<220>

<221> MISC\_FEATURE

<222> (2)..(2)

X-13991.ST25.txt

&lt;223&gt; Xaa at position 2 is Ala, Gly, Val, Leu, Ile, Ser, or Thr;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (6)..(6)

&lt;223&gt; Xaa at position 6 is Phe, Trp, or Tyr;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (10)..(10)

&lt;223&gt; Xaa at position 10 is Val, Trp, Ile, Leu, Phe, or Tyr;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (12)..(12)

&lt;223&gt; Xaa at position 12 is Ser, Trp, Tyr, Phe, Lys, Ile, Leu, Val;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (13)..(13)

&lt;223&gt; Xaa at position 13 is Tyr, Trp, or Phe;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (14)..(14)

&lt;223&gt; Xaa at position 14 is Leu, Phe, Tyr, or Trp;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (16)..(16)

&lt;223&gt; Xaa at position 16 is Gly, Glu, Asp, or Lys;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

X-13991.ST25.txt

&lt;222&gt; (19)..(19)

&lt;223&gt; Xaa at position 19 is Ala, Val, Ile, or Leu;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (21)..(21)

&lt;223&gt; Xaa at position 21 is Glu, Ile, or Ala;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (24)..(24)

&lt;223&gt; Xaa at position 24 is Ala or Glu;

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (27)..(27)

&lt;223&gt; Xaa at position 27 is Val or Ile; and

&lt;220&gt;

&lt;221&gt; MOD\_RES

&lt;222&gt; (30)..(30)

&lt;223&gt; AMIDATION

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (31)..(31)

&lt;223&gt; Xaa at position 31 is Gly, His or is absent.

&lt;400&gt; 12

Xaa	Xaa	Glu	Gly	Thr	Xaa	Thr	Ser	Asp	Xaa	Ser	Xaa	Xaa	Xaa	Glu	Xaa
1				5					10					15	

Gln	Ala	Xaa	Lys	Xaa	Phe	Ile	Xaa	Trp	Leu	Xaa	Lys	Gly	Arg	Xaa
			20					25					30	

&lt;210&gt; 13



X-13991.ST25.txt

&lt;211&gt; 616

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 13

His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Asp  
 20 25 30

Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu Glu  
 35 40 45

Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln Gln  
 50 55 60

Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu Phe  
 65 70 75 80

Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys Ser  
 85 90 95

Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu Arg  
 100 105 110

Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro Glu  
 115 120 125

Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu Pro  
 130 135 140

Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His Asp  
 145 150 155 160

Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg Arg  
 165 170 175

His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg Tyr  
 180 185 190

Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala Cys  
 195 200 205

Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser Ser  
 210 215 220

X-13991.ST25.txt

Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu Arg  
 225 230 235 240  
 Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro Lys  
 245 250 255  
 Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys Val  
 260 265 270  
 His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp Arg  
 275 280 285  
 Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser Ser  
 290 295 300  
 Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His Cys  
 305 310 315 320  
 Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser Leu  
 325 330 335  
 Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala Glu  
 340 345 350  
 Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg Arg  
 355 360 365  
 His Pro Asp Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys Thr Tyr  
 370 375 380  
 Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu Cys  
 385 390 395 400  
 Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu Pro Gln  
 405 410 415  
 Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly Glu Tyr  
 420 425 430  
 Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro Gln  
 435 440 445  
 Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys Val  
 450 455 460  
 Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys Ala  
 465 470 475 480  
 Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His Glu  
 485 490 495

X-13991.ST25.txt

Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser Leu  
500 505 510

Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr Tyr  
515 520 525

Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp Ile  
530 535 540

Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala Leu  
545 550 555 560

Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu Lys  
565 570 575

Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys Ala  
580 585 590

Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val Ala  
595 600 605

Ala Ser Gln Ala Ala Leu Gly Leu  
610 615

<210> 14

<211> 631

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 14

His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly  
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Gly  
20 25 30

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ala  
35 40 45

His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu Glu Asn  
50 55 60

Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln Gln Cys  
65 70 75 80

X-13991.ST25.txt

Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu Phe Ala  
 85 90 95  
 Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys Ser Leu  
 100 105 110  
 His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu Arg Glu  
 115 120 125  
 Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro Glu Arg  
 130 135 140  
 Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu Pro Arg  
 145 150 155 160  
 Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His Asp Asn  
 165 170 175  
 Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg Arg His  
 180 185 190  
 Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg Tyr Lys  
 195 200 205  
 Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala Cys Leu  
 210 215 220  
 Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser Ser Ala  
 225 230 235 240  
 Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu Arg Ala  
 245 250 255  
 Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro Lys Ala  
 260 265 270  
 Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys Val His  
 275 280 285  
 Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp Arg Ala  
 290 295 300  
 Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser Ser Lys  
 305 310 315 320  
 Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His Cys Ile  
 325 330 335  
 Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser Leu Ala  
 340 345 350

X-13991.ST25.txt

Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala Glu Ala  
355 360 365

Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg Arg His  
370 375 380

Pro Asp Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys Thr Tyr Glu  
385 390 395 400

Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu Cys Tyr  
405 410 415

Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu Pro Gln Asn  
420 425 430

Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly Glu Tyr Lys  
435 440 445

Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro Gln Val  
450 455 460

Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys Val Gly  
465 470 475 480

Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys Ala Glu  
485 490 495

Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His Glu Lys  
500 505 510

Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser Leu Val  
515 520 525

Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr Tyr Val  
530 535 540

Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp Ile Cys  
545 550 555 560

Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala Leu Val  
565 570 575

Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu Lys Ala  
580 585 590

Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys Ala Asp  
595 600 605

Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val Ala Ala  
610 615 620

X-13991.ST25.txt

Ser Gln Ala Ala Leu Gly Leu  
625 630

<210> 15

<211> 640

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 15

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
20 25 30

Ser Gly Ala Pro Pro Pro Ser Gly Gly Gly Gly Ser Gly Gly Gly  
35 40 45

Gly Ser Gly Gly Gly Gly Ser Asp Ala His Lys Ser Glu Val Ala His  
50 55 60

Arg Phe Lys Asp Leu Gly Glu Glu Asn Phe Lys Ala Leu Val Leu Ile  
65 70 75 80

Ala Phe Ala Gln Tyr Leu Gln Gln Cys Pro Phe Glu Asp His Val Lys  
85 90 95

Leu Val Asn Glu Val Thr Glu Phe Ala Lys Thr Cys Val Ala Asp Glu  
100 105 110

Ser Ala Glu Asn Cys Asp Lys Ser Leu His Thr Leu Phe Gly Asp Lys  
115 120 125

Leu Cys Thr Val Ala Thr Leu Arg Glu Thr Tyr Glu Met Ala Asp  
130 135 140

Cys Cys Ala Lys Gln Glu Pro Glu Arg Asn Glu Cys Phe Leu Gln His  
145 150 155 160

Lys Asp Asp Asn Pro Asn Leu Pro Arg Leu Val Arg Pro Glu Val  
165 170 175

Val Met Cys Thr Ala Phe His Asp Asn Glu Glu Thr Phe Leu Lys Lys  
180 185 190

X-13991.ST25.txt

Tyr Leu Tyr Glu Ile Ala Arg Arg His Pro Tyr Phe Tyr Ala Pro Glu  
 195 200  
 Leu Leu Phe Phe Ala Lys Arg Tyr Lys Ala Ala Phe Thr Glu Cys Cys  
 210 215  
 Gln Ala Ala Asp Lys Ala Ala Cys Leu Leu Pro Lys Leu Asp Glu Leu  
 225 230 235 240  
 Arg Asp Glu Gly Lys Ala Ser Ser Ala Lys Gln Arg Leu Lys Cys Ala  
 245 250 255  
 Ser Leu Gln Lys Phe Gly Glu Arg Ala Phe Lys Ala Trp Ala Val Ala  
 260 265 270  
 Arg Leu Ser Gln Arg Phe Pro Lys Ala Glu Phe Ala Glu Val Ser Lys  
 275 280 285  
 Leu Val Thr Asp Leu Thr Lys Val His Thr Glu Cys Cys His Gly Asp  
 290 295 300  
 Leu Leu Glu Cys Ala Asp Asp Arg Ala Asp Leu Ala Lys Tyr Ile Cys  
 305 310 315 320  
 Glu Asn Gln Asp Ser Ile Ser Ser Lys Leu Lys Glu Cys Cys Glu Lys  
 325 330 335  
 Pro Leu Leu Glu Lys Ser His Cys Ile Ala Glu Val Glu Asn Asp Glu  
 340 345 350  
 Met Pro Ala Asp Leu Pro Ser Leu Ala Ala Asp Phe Val Glu Ser Lys  
 355 360 365  
 Asp Val Cys Lys Asn Tyr Ala Glu Ala Lys Asp Val Phe Leu Gly Met  
 370 375 380  
 Phe Leu Tyr Glu Tyr Ala Arg Arg His Pro Asp Tyr Ser Val Val Leu  
 385 390 395 400  
 Leu Leu Arg Leu Ala Lys Thr Tyr Glu Thr Thr Leu Glu Lys Cys Cys  
 405 410 415  
 Ala Ala Ala Asp Pro His Glu Cys Tyr Ala Lys Val Phe Asp Glu Phe  
 420 425 430  
 Lys Pro Leu Val Glu Glu Pro Gln Asn Leu Ile Lys Gln Asn Cys Glu  
 435 440 445  
 Leu Phe Glu Gln Leu Gly Glu Tyr Lys Phe Gln Asn Ala Leu Leu Val  
 450 455 460

X-13991.ST25.txt

Arg Tyr Thr Lys Lys Val Pro Gln Val Ser Thr Pro Thr Leu Val Glu  
465 470 475 480

Val Ser Arg Asn Leu Gly Lys Val Gly Ser Lys Cys Cys Lys His Pro  
485 490 495

Glu Ala Lys Arg Met Pro Cys Ala Glu Asp Tyr Leu Ser Val Val Leu  
500 505 510

Asn Gln Leu Cys Val Leu His Glu Lys Thr Pro Val Ser Asp Arg Val  
515 520 525

Thr Lys Cys Cys Thr Glu Ser Leu Val Asn Arg Arg Pro Cys Phe Ser  
530 535 540

Ala Leu Glu Val Asp Glu Thr Tyr Val Pro Lys Glu Phe Asn Ala Glu  
545 550 555 560

Thr Phe Thr Phe His Ala Asp Ile Cys Thr Leu Ser Glu Lys Glu Arg  
565 570 575

Gln Ile Lys Lys Gln Thr Ala Leu Val Glu Leu Val Lys His Lys Pro  
580 585 590

Lys Ala Thr Lys Glu Gln Leu Lys Ala Val Met Asp Asp Phe Ala Ala  
595 600 605

Phe Val Glu Lys Cys Cys Lys Ala Asp Asp Lys Glu Thr Cys Phe Ala  
610 615 620

Glu Glu Gly Lys Lys Leu Val Ala Ala Ser Gln Ala Ala Leu Gly Leu  
625 630 635 640

<210> 16

<211> 624

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 16

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
1 5 10 15

Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser  
20 25 30



X-13991.ST25.txt

Ser Gly Ala Pro Pro Pro Ser Asp Ala His Lys Ser Glu Val Ala His  
 35 40 45  
 Arg Phe Lys Asp Leu Gly Glu Asn Phe Lys Ala Leu Val Leu Ile  
 50 55 60  
 Ala Phe Ala Gln Tyr Leu Gln Gln Cys Pro Phe Glu Asp His Val Lys  
 65 70 75 80  
 Leu Val Asn Glu Val Thr Glu Phe Ala Lys Thr Cys Val Ala Asp Glu  
 85 90 95  
 Ser Ala Glu Asn Cys Asp Lys Ser Leu His Thr Leu Phe Gly Asp Lys  
 100 105 110  
 Leu Cys Thr Val Ala Thr Leu Arg Glu Thr Tyr Gly Glu Met Ala Asp  
 115 120 125  
 Cys Cys Ala Lys Gln Glu Pro Glu Arg Asn Glu Cys Phe Leu Gln His  
 130 135 140  
 Lys Asp Asp Asn Pro Asn Leu Pro Arg Leu Val Arg Pro Glu Val Asp  
 145 150 155 160  
 Val Met Cys Thr Ala Phe His Asp Asn Glu Glu Thr Phe Leu Lys Lys  
 165 170 175  
 Tyr Leu Tyr Glu Ile Ala Arg Arg His Pro Tyr Phe Tyr Ala Pro Glu  
 180 185 190  
 Leu Leu Phe Phe Ala Lys Arg Tyr Lys Ala Ala Phe Thr Glu Cys Cys  
 195 200 205  
 Gln Ala Ala Asp Lys Ala Ala Cys Leu Leu Pro Lys Leu Asp Glu Leu  
 210 215 220  
 Arg Asp Glu Gly Lys Ala Ser Ser Ala Lys Gln Arg Leu Lys Cys Ala  
 225 230 235 240  
 Ser Leu Gln Lys Phe Gly Glu Arg Ala Phe Lys Ala Trp Ala Val Ala  
 245 250 255  
 Arg Leu Ser Gln Arg Phe Pro Lys Ala Glu Phe Ala Glu Val Ser Lys  
 260 265 270  
 Leu Val Thr Asp Leu Thr Lys Val His Thr Glu Cys Cys His Gly Asp  
 275 280 285  
 Leu Leu Glu Cys Ala Asp Asp Arg Ala Asp Leu Ala Lys Tyr Ile Cys  
 290 295 300  
 35

X-13991.ST25.txt

Glu Asn Gln Asp Ser Ile Ser Ser Lys Leu Lys Glu Cys Cys Glu Lys  
 305 310 315 320  
 Pro Leu Leu Glu Lys Ser His Cys Ile Ala Glu Val Glu Asn Asp Glu  
 325 330 335  
 Met Pro Ala Asp Leu Pro Ser Leu Ala Ala Asp Phe Val Glu Ser Lys  
 340 345 350  
 Asp Val Cys Lys Asn Tyr Ala Glu Ala Lys Asp Val Phe Leu Gly Met  
 355 360 365  
 Phe Leu Tyr Glu Tyr Ala Arg Arg His Pro Asp Tyr Ser Val Val Leu  
 370 375 380  
 Leu Leu Arg Leu Ala Lys Thr Tyr Glu Thr Thr Leu Glu Lys Cys Cys  
 385 390 395 400  
 Ala Ala Ala Asp Pro His Glu Cys Tyr Ala Lys Val Phe Asp Glu Phe  
 405 410 415  
 Lys Pro Leu Val Glu Glu Pro Gln Asn Leu Ile Lys Gln Asn Cys Glu  
 420 425 430  
 Leu Phe Glu Gln Leu Gly Glu Tyr Lys Phe Gln Asn Ala Leu Leu Val  
 435 440 445  
 Arg Tyr Thr Lys Lys Val Pro Gln Val Ser Thr Pro Thr Leu Val Glu  
 450 455 460  
 Val Ser Arg Asn Leu Gly Lys Val Gly Ser Lys Cys Cys Lys His Pro  
 465 470 475 480  
 Glu Ala Lys Arg Met Pro Cys Ala Glu Asp Tyr Leu Ser Val Val Leu  
 485 490 495  
 Asn Gln Leu Cys Val Leu His Glu Lys Thr Pro Val Ser Asp Arg Val  
 500 505 510  
 Thr Lys Cys Cys Thr Glu Ser Leu Val Asn Arg Arg Pro Cys Phe Ser  
 515 520 525  
 Ala Leu Glu Val Asp Glu Thr Tyr Val Pro Lys Glu Phe Asn Ala Glu  
 530 535 540  
 Thr Phe Thr Phe His Ala Asp Ile Cys Thr Leu Ser Glu Lys Glu Arg  
 545 550 555 560  
 Gln Ile Lys Lys Gln Thr Ala Leu Val Glu Leu Val Lys His Lys Pro  
 565 570 575

X-13991.ST25.txt

Lys Ala Thr Lys Glu Gln Leu Lys Ala Val Met Asp Asp Phe Ala Ala  
580 585 590

Phe Val Glu Lys Cys Cys Lys Ala Asp Asp Lys Glu Thr Cys Phe Ala  
595 600 605

Glu Glu Gly Lys Lys Leu Val Ala Ala Ser Gln Ala Ala Leu Gly Leu  
610 615 620

<210> 17

<211> 640

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 17

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
1 5 10 15

Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser  
20 25 30

Ser Gly Ala Pro Pro Pro Ser Gly Gly Gly Gly Gly Ser Gly Gly Gly  
35 40 45

Gly Ser Gly Gly Gly Gly Ser Asp Ala His Lys Ser Glu Val Ala His  
50 55 60

Arg Phe Lys Asp Leu Gly Glu Glu Asn Phe Lys Ala Leu Val Leu Ile  
65 70 75 80

Ala Phe Ala Gln Tyr Leu Gln Gln Cys Pro Phe Glu Asp His Val Lys  
85 90 95

Leu Val Asn Glu Val Thr Glu Phe Ala Lys Thr Cys Val Ala Asp Glu  
100 105 110

Ser Ala Glu Asn Cys Asp Lys Ser Leu His Thr Leu Phe Gly Asp Lys  
115 120 125

Leu Cys Thr Val Ala Thr Leu Arg Glu Thr Tyr Gly Glu Met Ala Asp  
130 135 140

Cys Cys Ala Lys Gln Glu Pro Glu Arg Asn Glu Cys Phe Leu Gln His  
145 150 155 160

X-13991.ST25.txt

Lys Asp Asp Asn Pro Asn Leu Pro Arg Leu Val Arg Pro Glu Val Asp  
 165 170 175  
 Val Met Cys Thr Ala Phe His Asp Asn Glu Glu Thr Phe Leu Lys Lys  
 180 185 190  
 Tyr Leu Tyr Glu Ile Ala Arg Arg His Pro Tyr Phe Tyr Ala Pro Glu  
 195 200 205  
 Leu Leu Phe Phe Ala Lys Arg Tyr Lys Ala Ala Phe Thr Glu Cys Cys  
 210 215 220  
 Gln Ala Ala Asp Lys Ala Ala Cys Leu Leu Pro Lys Leu Asp Glu Leu  
 225 230 235 240  
 Arg Asp Glu Gly Lys Ala Ser Ser Ala Lys Gln Arg Leu Lys Cys Ala  
 245 250 255  
 Ser Leu Gln Lys Phe Gly Glu Arg Ala Phe Lys Ala Trp Ala Val Ala  
 260 265 270  
 Arg Leu Ser Gln Arg Phe Pro Lys Ala Glu Phe Ala Glu Val Ser Lys  
 275 280 285  
 Leu Val Thr Asp Leu Thr Lys Val His Thr Glu Cys Cys His Gly Asp  
 290 295 300  
 Leu Leu Glu Cys Ala Asp Asp Arg Ala Asp Leu Ala Lys Tyr Ile Cys  
 305 310 315 320  
 Glu Asn Gln Asp Ser Ile Ser Ser Lys Leu Lys Glu Cys Cys Glu Lys  
 325 330 335  
 Pro Leu Leu Glu Lys Ser His Cys Ile Ala Glu Val Glu Asn Asp Glu  
 340 345 350  
 Met Pro Ala Asp Leu Pro Ser Leu Ala Ala Asp Phe Val Glu Ser Lys  
 355 360 365  
 Asp Val Cys Lys Asn Tyr Ala Glu Ala Lys Asp Val Phe Leu Gly Met  
 370 375 380  
 Phe Leu Tyr Glu Tyr Ala Arg Arg His Pro Asp Tyr Ser Val Val Leu  
 385 390 395 400  
 Leu Leu Arg Leu Ala Lys Thr Tyr Glu Thr Thr Leu Glu Lys Cys Cys  
 405 410 415  
 Ala Ala Ala Asp Pro His Glu Cys Tyr Ala Lys Val Phe Asp Glu Phe  
 420 425 430

X-13991.ST25.txt

Lys Pro Leu Val Glu Glu Pro Gln Asn Leu Ile Lys Gln Asn Cys Glu  
 435 440  
 Leu Phe Glu Gln Leu Gly Glu Tyr Lys Phe Gln Asn Ala Leu Leu Val  
 450 455 460  
 Arg Tyr Thr Lys Lys Val Pro Gln Val Ser Thr Pro Thr Leu Val Glu  
 465 470 475 480  
 Val Ser Arg Asn Leu Gly Lys Val Gly Ser Lys Cys Cys Lys His Pro  
 485 490 495  
 Glu Ala Lys Arg Met Pro Cys Ala Glu Asp Tyr Leu Ser Val Val Leu  
 500 505 510  
 Asn Gln Leu Cys Val Leu His Glu Lys Thr Pro Val Ser Asp Arg Val  
 515 520 525  
 Thr Lys Cys Cys Thr Glu Ser Leu Val Asn Arg Arg Pro Cys Phe Ser  
 530 535 540  
 Ala Leu Glu Val Asp Glu Thr Tyr Val Pro Lys Glu Phe Asn Ala Glu  
 545 550 555 560  
 Thr Phe Thr Phe His Ala Asp Ile Cys Thr Leu Ser Glu Lys Glu Arg  
 565 570 575  
 Gln Ile Lys Lys Gln Thr Ala Leu Val Glu Leu Val Lys His Lys Pro  
 580 585 590  
 Lys Ala Thr Lys Glu Gln Leu Lys Ala Val Met Asp Asp Phe Ala Ala  
 595 600 605  
 Phe Val Glu Lys Cys Cys Lys Ala Asp Asp Lys Glu Thr Cys Phe Ala  
 610 615 620  
 Glu Glu Gly Lys Lys Leu Val Ala Ala Ser Gln Ala Ala Leu Gly Leu  
 625 630 635 640

&lt;210&gt; 18

&lt;211&gt; 264

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

X-13991.ST25.txt

&lt;400&gt; 18

His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ala  
 20 25 30

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala  
 35 40 45

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro  
 50 55 60

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val  
 65 70 75 80

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val  
 85 90 95

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln  
 100 105 110

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln  
 115 120 125

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala  
 130 135 140

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro  
 145 150 155 160

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr  
 165 170 175

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser  
 180 185 190

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr  
 195 200 205

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr  
 210 215 220

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe  
 225 230 235 240

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys  
 245 250 255

Ser Leu Ser Leu Ser Pro Gly Lys  
 260

X-13991.ST25.txt

&lt;210&gt; 19

&lt;211&gt; 272

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 19

His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
 20 25 30

Ser Gly Ala Pro Pro Pro Ser Ala Glu Pro Lys Ser Cys Asp Lys Thr  
 35 40 45

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
 50 55 60

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
 65 70 75 80

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
 85 90 95

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
 100 105 110

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
 115 120 125

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
 130 135 140

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
 145 150 155 160

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
 165 170 175

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
 180 185 190

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
 195 200 205

X-13991.ST25.txt

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
 210 215

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
 225 230 235 240

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
 245 250 255

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 260 265 270

<210> 20

<211> 264

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 20

His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ala  
 20 25 30

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala  
 35 40 45

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro  
 50 55 60

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val  
 65 70 75 80

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val  
 85 90 95

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln  
 100 105 110

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln  
 115 120

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala  
 130 135 140



X-13991.ST25.txt

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro  
145 150 155 160

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr  
165 170 175

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser  
180 185 190

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr  
195 200 205

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr  
210 215 220

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe  
225 230 235 240

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys  
245 250 255

Ser Leu Ser Leu Ser Pro Gly Lys  
260

<210> 21

<211> 272

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 21

His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
20 25 30

Ser Gly Ala Pro Pro Pro Ser Ala Glu Pro Lys Ser Cys Asp Lys Thr  
35 40 45

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
50 55 60

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
65 70 75 80

X-13991.ST25.txt

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
85 95

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
100 105 110

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
115 120 125

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
130 135 140

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
145 150 155 160

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
165 170 175

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
180 185 190

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
195 200 205

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
210 215 220

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
225 230 235 240

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
245 250 255

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
260 265 270

&lt;210&gt; 22

&lt;211&gt; 272

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 22

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
1 5 10 15

X-13991.ST25.txt

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
 20 25 30  
 Ser Gly Ala Ser Ser Gly Ala Ala Glu Pro Lys Ser Cys Asp Lys Thr  
 35 40 45  
 His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
 50 55 60  
 Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
 65 70 75 80  
 Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
 85 90 95  
 Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
 100 105 110  
 Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
 115 120 125  
 Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
 130 135 140  
 Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
 145 150 155 160  
 Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
 165 170 175  
 Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
 180 185 190  
 Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
 195 200 205  
 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
 210 215 220  
 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
 225 230 235 240  
 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
 245 250 255  
 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 260 265 270

&lt;210&gt; 23

X-13991.ST25.txt

&lt;211&gt; 287

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 23

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
 20 25 30

Ser Gly Ala Pro Pro Pro Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly  
 35 40 45

Ser Gly Gly Gly Gly Ser Ala Glu Pro Lys Ser Cys Asp Lys Thr His  
 50 55 60

Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val  
 65 70 75 80

Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr  
 85 90 95

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu  
 100 105 110

Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys  
 115 120 125

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser  
 130 135 140

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys  
 145 150 155 160

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile  
 165 170 175

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro  
 180 185 190

Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu  
 195 200 205

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn  
 210 215 220

X-13991.ST25.txt

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser  
 225 230 235 240

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg  
 245 250 255

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu  
 260 265 270

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 275 280 285

<210> 24

<211> 284

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 24

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
 20 25 30

Ser Gly Ala Pro Pro Pro Ser Gly Gly Gly Ser Gly Gly Gly Gly  
 35 40 45

Ser Gly Gly Gly Ser Ala Glu Ser Lys Tyr Gly Pro Pro Cys Pro  
 50 55 60

Ser Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe  
 65 70 75 80

Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val  
 85 90 95

Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe  
 100 105 110

Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro  
 115 120 125

Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr  
 130 135 140

X-13991.ST25.txt

Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val  
 145 150 155 160

Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala  
 165 170 175

Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln  
 180 185 190

Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly  
 195 200 205

Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro  
 210 215 220

Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser  
 225 230 235 240

Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu  
 245 250 255

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His  
 260 265 270

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys  
 275 280

&lt;210&gt; 25

&lt;211&gt; 302

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 25

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
 20 25 30

Ser Gly Ala Pro Pro Pro Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly  
 35 40 45

Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser  
 50 55 60

X-13991.ST25.txt

Gly Gly Gly Gly Ser Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr  
65 70 75 80

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe  
85 90 95

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro  
100 105 110

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val  
115 120 125

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr  
130 135 140

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val  
145 150 155 160

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys  
165 170 175

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser  
180 185 190

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro  
195 200 205

Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val  
210 215 220

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly  
225 230 235 240

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Val Leu Asp Ser Asp  
245 250 255

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp  
260 265 270

Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His  
275 280 285

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
290 295 300

<210> 26

<211> 294

<212> PRT

X-13991.ST25.txt

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 26

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Gly  
 20 25 30

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly  
 35 40 45

Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ala Glu Pro  
 50 55 60

Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu  
 65 70 75 80

Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp  
 85 90 95

Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp  
 100 105 110

Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly  
 115 120 125

Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn  
 130 135 140

Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp  
 145 150 155 160

Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro  
 165 170 175

Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu  
 180 185 190

Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn  
 195 200 205

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile  
 210 215 220

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr  
 225 230 235 240



X-13991.ST25.txt

Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys  
245 250 255

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys  
260 265 270

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu  
275 280 285

Ser Leu Ser Pro Gly Lys  
290

<210> 27

<211> 280

<212> PRT

<213> Artificial sequence

<220>

<223> synthetic construct

<400> 27

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly Ser  
20 25 30

Ser Gly Ala Pro Pro Pro Ser Ser Ser Gly Ala Pro Pro Pro Ser Ala  
35 40 45

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala  
50 55 60

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro  
65 70 75 80

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val  
85 90 95

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val  
100 105 110

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln  
115 120 125

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln  
130 135 140

X-13991.ST25.txt

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala  
 145 150 155 160

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro  
 165 170 175

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr  
 180 185 190

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser  
 195 200 205

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr  
 210 215 220

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr  
 225 230 235 240

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe  
 245 250 255

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys  
 260 265 270

Ser Leu Ser Leu Ser Pro Gly Lys  
 275 280

&lt;210&gt; 28

&lt;211&gt; 287

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 28

His Gly Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Glu  
 1 5 10 15

Gln Ala Val Lys Glu Phe Ile Ala Trp Leu Ile Lys Gly Arg Gly Ser  
 20 25 30

Ser Gly Ala Pro Pro Pro Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly  
 35 40 45

Ser Gly Gly Gly Gly Ser Ala Glu Pro Lys Ser Cys Asp Lys Thr His  
 50 55 60

X-13991.ST25.txt

Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val  
65 70 75 80

Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr  
85 90 95

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu  
100 105 110

Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys  
115 120 125

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser  
130 135 140

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys  
145 150 155 160

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile  
165 170 175

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro  
180 185 190

Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu  
195 200 205

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn  
210 215 220

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser  
225 230 235 240

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg  
245 250 255

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu  
260 265 270

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
275 280 285

&lt;210&gt; 29

&lt;211&gt; 272

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

X-13991.ST25.txt

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 29

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
 1 5 10 15

Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser  
 20 25 30

Ser Gly Ala Pro Pro Pro Ser Ala Glu Pro Lys Ser Cys Asp Lys Thr  
 35 40 45

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
 50 55 60

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
 65 70 75 80

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
 85 90 95

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
 100 105 110

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
 115 120 125

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
 130 135 140

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
 145 150 155 160

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
 165 170 175

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
 180 185 190

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
 195 200 205

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
 210 215 220

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
 225 230 235 240

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
 245 250 255

X-13991.ST25.txt

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 260 265 270

&lt;210&gt; 30

&lt;211&gt; 272

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; synthetic construct

&lt;400&gt; 30

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
 1 5 10 15

Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser  
 20 25 30

Ser Gly Ala Ser Ser Gly Ala Ala Glu Pro Lys Ser Cys Asp Lys Thr  
 35 40 45

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser  
 50 55 60

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg  
 65 70 75 80

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro  
 85 90 95

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala  
 100 105 110

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val  
 115 120 125

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr  
 130 135 140

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr  
 145 150 155 160

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
 165 170 175

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys  
 180 185 190

X-13991.ST25.txt

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
195 200

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
210 215

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
225 230 235 240

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
245 250 255

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
260 265 270

<210> 31

<211> 287

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 31

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
1 5 10 15

Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser  
20 25 30

Ser Gly Ala Pro Pro Pro Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly  
35 40 45

Ser Gly Gly Gly Gly Ser Ala Glu Pro Lys Ser Cys Asp Lys Thr His  
50 55 60

Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val  
65 70 75 80

Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr  
85 90 95

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu  
100 105 110

Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys  
115 120 125

X-13991.ST25.txt

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser  
 130 135 140

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys  
 145 150 155 160

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile  
 165 170 175

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro  
 180 185 190

Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu  
 195 200 205

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn  
 210 215 220

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser  
 225 230 235 240

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg  
 245 250 255

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu  
 260 265 270

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 275 280 285

<210> 32

<211> 232

<212> PRT

<213> Homo sapiens

<400> 32.

Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro  
 1 5 10 15

Ala Pro Glu Lys Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro  
 20 25 30

Lys Asp Thr Lys Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val  
 35 40 45

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val  
 50 55 60

X-13991.ST25.txt

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln  
65 70 75 80

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln  
85 90 95

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala  
100 105 110

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro  
115 120 125

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr  
130 135 140

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser  
145 150 155 160

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr  
165 170 175

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr  
180 185 190

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe  
195 200 205

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys  
210 215 220

Ser Leu Ser Leu Ser Pro Gly Lys  
225 230

&lt;210&gt; 33

&lt;211&gt; 703

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

<400> 33  
gagcccaaat cttgtgacaa aactcacaca tgcccaccgt gccccagcacc tgaactcctg 60  
gggggaccgt cagtcttctt cttccccca aaaccaagg acaccctcat gatctcccg 120  
acccttgagg tcacatgcgt ggtggtggac gtgagccacg aagaccctga ggtcaagttc 180  
aactggtacg tggacggcgt ggaggtgcat aatgccaga caaagccgag ggaggagcag 240  
tacaacagca cgtaccgtgt ggtcagcgtc ctcaccgtcc tgcaccagga ctggctgaat 300  
ggcaaggagt acaagtgcga ggtctccaac aaagccctcc cagcccccat cgagaaaacc 360



X-13991.ST25.txt

```

atctccaag ccaaggga gccccgagaa ccacaggtgt acacctgcc cccatcccgg 420
gaggagatga ccaagaacca ggtcagcctg acctgcctgg tcaaggctt ctatcccagc 480
gacatcgccg tggagtggga gagcaatggg cagccggaga acaactacaa gaccacgctt 540
cccgtagctg actccgacgg ctctctcttc ctctatagca agctcaccgt ggacaagagc 600
agggtgcagc aggggaacgt cttctcatgc tccgtgatgc atgaggctct gcacaaccac 660
tacacgcaga agagcctctc cctgtctccg ggtaaatgat agt 703

```

&lt;210&gt; 34

&lt;211&gt; 585

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

```

Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu
1      5      10      15

```

```

Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln
20      25      30

```

```

Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu
35      40      45

```

```

Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys
50      55      60

```

```

Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu
65      70      75      80

```

```

Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro
85      90      95

```

```

Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu
100     105     110

```

```

Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His
115     120     125

```

```

Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg
130     135     140

```

```

Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg
145     150     155     160

```

```

Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala
165     170     175

```

X-13991.ST25.txt

Cys Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser  
 180 185 190  
 Ser Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu  
 195 200 205  
 Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro  
 210 215 220  
 Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys  
 225 230 235 240  
 Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp  
 245 250 255  
 Arg Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser  
 260 265 270  
 Ser Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His  
 275 280 285  
 Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser  
 290 295 300  
 Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala  
 305 310 315 320  
 Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg  
 325 330 335  
 Arg His Pro Asp Tyr Ser Val Val Leu Leu Arg Leu Ala Lys Thr  
 340 345 350  
 Tyr Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu  
 355 360 365  
 Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu Pro  
 370 375 380  
 Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Asn Leu Gly Glu  
 385 390 395 400  
 Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro  
 405 410 415  
 Gln Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys  
 420 425 430  
 Val Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys  
 435 440 445

X-13991.ST25.txt

Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His  
450 455 460

Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser  
465 470 475 480

Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr  
485 490 495

Tyr Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp  
500 505 510

Ile Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala  
515 520 525

Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu  
530 535 540

Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys  
545 550 555 560

Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val  
565 570 575

Ala Ala Ser Gln Ala Ala Leu Gly Leu  
580 585

&lt;210&gt; 35

&lt;211&gt; 1762

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

<400> 35  
gatgcgcaca agagtggagt tgctcatcgg tttaagatt tgggagaaga aaatttcaaa 60  
gccttgggtg tgattgcctt tgctcagtat cttcagcagt gtccatttga agatcatgta 120  
aaattagtga atgaagtaac tgaatttgca aaaacatgtg ttctgatga gtcagctgaa 180  
aattgtgaca aatcacttca tacccttttt ggagacaaat tatgcacagt tgcaactctt 240  
cgtgaaacct atggtgaaat ggcgtgactgc tgtgcaaaac aagaacctga gagaaatgaa 300  
tgcttcttgc aacacaaaga tgacaaccca aacctcccc gattggtgag accagaggtt 360  
gatgtgatgt gcactgcttt tcatgacaat gaagagacat ttttgaaaa atacttatat 420  
gaaattgcca gaagacatcc ttacttttat gcccgggaac tccttttctt tgcataaagg 480  
tataaagctg cttttacaga atgttgccaa gctgctgata aagctgcctg cctgttgcca 540  
aagctcgatg aacttcggga tgaaggggaag gcttcgtctg ccaaacagag actcaagtgt 600  
i 51

## x-13991.ST25.txt

gccagctctcc	aaaaatttgg	agaaagagct	ttcaaagcat	gggcagtagc	tcgcctgagc	660
cagagatttc	ccaagctga	gtttgcagaa	gtttccaagt	tagtgacaga	tcttaccaaa	720
gtccacacgg	aatgctgcca	tggagatctg	cttgaatgtg	ctgatgacag	ggcggacctt	780
gccaaagtata	tctgtgaaaa	tcaagattcg	atctccagta	aactgaagga	atgctgtgaa	840
aaacctctgt	tggaaaaaat	ccactgcatt	gccgaagtgg	aaaatgatga	gatgcctgct	900
gacttgcctt	cattagctgc	tgattttgtt	gaaagtaagg	atgtttgcaa	aaactatgct	960
gaggcaaagg	atgtcttcct	gggcatgttt	ttgtatgaat	atgcaagaag	gcctcctgat	1020
tactctgtcg	tgctgtgctg	gagacttgcc	aagacatatg	aaaccactct	agagaagtgct	1080
tggtccgctg	cagatcctca	tgaatgctat	gccaaagtgt	tcgatgaatt	taaacctctt	1140
gtggaagagc	ctcagaattt	aatcaacaaa	aattgtgagc	tttttgagca	gcttgagagag	1200
tacaaatttc	agaatgcgct	attagtctgt	tacaccaaga	aagtacccca	agtgctcaact	1260
ccaactcttg	tagaggctc	aagaaaccta	ggaaaagtgg	gcagcaaatg	ttgtaaacat	1320
cctgaagcaa	aaagaatgcc	ctgtgcagaa	gactatctat	ccgtggtcct	gaaccagtta	1380
tggtgtgttc	atgagaaaac	gccagtaagt	gacagagtca	ccaatgctg	cacagaatcc	1440
ttggtgaaca	ggcgacctg	cttttcagct	ctggaagtgc	atgaaacata	cgttcccaaa	1500
gagtttaatg	ctgaaacatt	caccttccat	gcagatatat	gcacacttcc	tgagaaggag	1560
agacaaatca	agaaaacaac	tgacttgttt	gagctcgtga	aacacaagcc	caaggcaaca	1620
aaagagcaac	tgaagctgt	tatggatgat	ttcgagctt	ttgtagagaa	gtgctgcaag	1680
gctgacgata	aggagacctg	ctttgccgag	gagggtaaaa	aacttgttgc	tgcaagtcaa	1740
gctgccttag	gcttataatg	ac				1762